

Article

In Vitro Anti-Leishmanial Activity of Essential Oils Extracted from Vietnamese Plants

Thanh Binh Le ^{1,2,*} , Claire Beaufay ¹ , Duc Trong Nghiem ³ ,
Marie-Paule Mingeot-Leclercq ⁴ and Joëlle Quetin-Leclercq ^{1,*}

¹ GNOS Research Group, Louvain Drug Research Institute, Université Catholique de Louvain, Bruxelles 1200, Belgium; claire.beaufay@uclouvain.be

² Department of Pharmacognosy, Hanoi University of Pharmacy, 13-15 Le Thanh Tong, Hoan Kiem, Hanoi 100000, Vietnam

³ Department of Botany, Hanoi University of Pharmacy, 13-15 Le Thanh Tong, Hoan Kiem, Hanoi 100000, Vietnam; ductrongeb@hup.edu.vn

⁴ TFAR Research Group, Louvain Drug Research Institute, Université Catholique de Louvain, Bruxelles 1200, Belgium; marie-paule.mingeot@uclouvain.be

* Correspondence: thanh.le@student.uclouvain.be (T.B.L.); joelle.leclercq@uclouvain.be (J.Q.-L.);
Tel.: +32-2-764-7234 (T.B.L.); +32-2-764-7254 (J.Q.-L.)

Received: 4 May 2017; Accepted: 22 June 2017; Published: 27 June 2017

Abstract: *Leishmania mexicana* is one of the pathogens causing cutaneous leishmaniasis which is associated with patient morbidity. In our researches for new safe and effective treatments, thirty-seven essential oils (EOs) extracted from Vietnamese plants were screened in vitro for the first time on *Leishmania mexicana mexicana* (*Lmm*) promastigotes at the maximum concentration of 50 nL/mL. Active EOs were also analyzed for cytotoxicity on mammalian cell lines (WI38, J774) and their selectivity indices (SI) were calculated. Their composition was determined by GC-MS and GC-FID. Our results indicated that EOs extracted from *Cinnamomum cassia*, *Zingiber zerumbet*, *Elsholtzia ciliata* and *Amomum aromaticum*, possessed a moderate anti-leishmanial activity, with IC₅₀ values of 2.92 ± 0.08, 3.34 ± 0.34, 8.49 ± 0.32 and 9.25 ± 0.64 nL/mL respectively. However, they also showed cytotoxicity with SI < 10. The most promising EO was extracted from *Ocimum gratissimum*, displaying an IC₅₀ of 4.85 ± 1.65 nL/mL and SI > 10. It contained 86.5% eugenol, which was demonstrated to be effective on *Lmm* with IC₅₀ of 2.57 ± 0.57 nL/mL and not toxic on mammalian cells, explaining the observed activity.

Keywords: *Leishmania mexicana mexicana*; essential oils; *Ocimum gratissimum*; *Cinnamomum cassia*; *Zingiber zerumbet*; *Elsholtzia ciliata*; *Amomum aromaticum*; eugenol

1. Introduction

Cutaneous leishmaniasis (CL), the most common form of leishmaniasis, largely affects poor and developing countries in Africa, the Mediterranean Basin, the Middle East, central Asia (the Old World) and South America (the New World). The estimated number of new CL cases worldwide ranges from 0.7 million to 1.3 million annually [1].

CL is caused by protozoan parasites of the *Leishmania* (*L.*) genus. Infected female phlebotomine sand flies inject these parasites from their proboscis to human when they take their blood meals. Skin lesions, the clinical signs of CL, develop within several weeks or months after exposure. These lesions are sometimes self-healing without treatment, or become chronic. In cases of ulcers healing, they leave permanent deep scars, which often cause a serious social prejudice. Conversely, if the disease becomes chronic, there is a high risk of severe bacterial infection. CL in the New World, mainly caused by *L. mexicana*, *L. amazonensis*, *L. venezuelensis*, *L. braziliensis*, *L. guyanensis*, *L. panamensis*, and *L. peruviana*,

is usually more severe and lasts longer than that in the Old World, which is caused by *L. tropica*, *L. major*, *L. aethiopica*, *L. infantum* and *L. donovani*. Moreover, CL due to *L. mexicana*, *L. amazonensis* and *L. panamensis* can develop to diffuse forms. Lesions can spread from the skin to the mucosal surfaces of the nose or the mouth, leading to chronic stuffiness, bleeding, and inflamed mucosa or sores. In advanced cases, it can lead to the ulcerative destruction of the nose (such as perforation of the nasal septum), mouth, and pharynx [1,2].

Drugs are the only therapeutic option for the treatment of CL as there is no vaccine at the moment. However, existing drugs, such as pentavalent antimonials and miltefosine, have serious drawbacks in terms of safety, efficacy, cost and difficulty in administration [3,4]. Moreover, diffuse CL initially responding to the standard treatment relapses and becomes unresponsive to further treatments [2]. Research on a safer, more effective, and shorter-course treatment for CL is therefore urgent. Nevertheless, the development of new medicines for treatment of this disease is challenging because of the variety of parasite species, pathology and immune responses [2].

Essential oils (EOs), also known as volatile oils, are natural products formed by a mixture of volatile compounds produced by many plants. They are known to be biologically active, mainly possessing antibacterial, antifungal, and antioxidant properties. There are more and more studies on EOs biological activity because they are usually devoid of long-term genotoxic risks [5]. Interestingly, topical application, the most frequent way to administer EOs, is recommended to be explored for the development of new anti-leishmanial drugs targeting CL because of their effectiveness and safety [2]. Several EOs showed interesting potential to be used as new anti-parasitic drugs such as those extracted from *Chenopodium ambrosioides* L. (a synonym of *Dysphania ambrosioides* (L.) Mosyakin & Clemant) [6,7] or *Bixa orellana* L. [8]. However, EOs are complex mixtures of several compounds and their compositions vary according to many factors, such as plant's environment and growing conditions, methods of harvesting, extraction and storage. Moreover, the major component of an EO can also vary in different chemotypes of the same plant species. This chemical variability can influence their activities or adverse effects. Therefore, a clear knowledge of the EO composition is necessary [9].

With the aim of discovering new natural products against *Leishmania mexicana mexicana* and developing active EOs or compounds for topical application, 37 Vietnamese plants were selected for anti-leishmanial investigation in our work. It is the first time that these Vietnamese EOs were tested on this *Leishmania* species. Moreover, this study also presents the results of chemical composition analyses of the most interesting samples.

2. Results

A primary screening was performed at concentrations of 50 and 25 nL/mL for the 37 EOs. The results showed that among them five EOs, extracted from *Amomum aromaticum*, *Cinnamomum cassia*, *Elsholtzia ciliata*, *Ocimum gratissimum*, and *Zingiber zerumbet*, displayed a promising effect against *Leishmania mexicana mexicana* (*Lmm*) promastigotes with less than 1% viable parasites at the lower concentration tested (Table 1).

Table 1. Viability of *Lmm* promastigotes in the presence of 25 nL/mL EO.

Plant Species	EO Obtained From	Viability of <i>Lmm</i> promastigotes (%) (Average \pm Standard Deviation)
<i>Ageratum conyzoides</i> (L.) L.	leaves	89.84 \pm 1.58
<i>Alpinia galanga</i> (L.) Willd.	rhizomes	93.02 \pm 4.90
<i>Amomum aromaticum</i> Roxb.	fruits	0.27 \pm 0.05
<i>Anomum schmidtii</i> (K. Schum.) Gagnep.	rhizomes	96.82 \pm 1.13
<i>Anethum graveolens</i> L.	fruits	96.53 \pm 1.26
<i>Artemisia annua</i> L.	leaves	92.32 \pm 1.55
<i>Blumea lanceolaria</i> (Roxb.) Druce	leaves	93.59 \pm 1.24
<i>Cinnamomum cassia</i> (L.) J. Presl	stem barks	0.48 \pm 0.01
<i>Clausena indica</i> (Dalzell) Oliv.	leaves	95.07 \pm 7.40
<i>Coriandrum sativum</i> L.	fruits	94.47 \pm 2.38

Table 1. Cont.

Plant Species	EO Obtained From	Viability of <i>Lmm</i> promastigotes (%) (Average \pm Standard Deviation)
<i>Curcuma longa</i> L.	rhizomes	92.90 \pm 1.61
<i>Curcuma zedoaria</i> (Christm.) Roscoe	rhizomes	99.53 \pm 0.23
<i>Dysphania ambrosioides</i> (L.) Mosyakin & Clemant	leaves, fruits	92.69 \pm 1.93
<i>Elsholtzia blanda</i> (Benth.) Benth.	leaves	94.70 \pm 6.12
<i>Elsholtzia ciliata</i> (Thunb.) Hyl.	leaves	0.38 \pm 0.00
<i>Elsholtzia communis</i> (Collett & Hemsl.) Diels	leaves	97.24 \pm 2.19
<i>Elsholtzia penduliflora</i> W. W. Sm.	leaves	97.88 \pm 0.37
<i>Eucalyptus camaldulensis</i> Dehnh.	leaves	101.13 \pm 0.54
<i>Hedychiium coronarium</i> J.Koenig	rhizomes	91.79 \pm 4.60
<i>Hyptis suaveolens</i> (L.) Poit.	leaves	67.32 \pm 1.97
<i>Illicium verum</i> Hook. f.	fruits	93.35 \pm 5.29
<i>Kaempferia galanga</i> L.	rhizomes	100.00 \pm 7.47
<i>Litsea cubeba</i> (Lour.) Pers.	fruits	96.00 \pm 5.18
<i>Litsea cubeba</i> (Lour.) Pers.	leaves	94.67 \pm 2.99
<i>Melaleuca alternifolia</i> (Maiden & Betch) Cheel	leaves	87.61 \pm 4.93
<i>Melaleuca cajuputi</i> Powell	leaves	92.53 \pm 3.17
<i>Ocimum gratissimum</i> L.	leaves	0.13 \pm 0.01
<i>Ocimum tenuiflorum</i> L.	leaves	89.08 \pm 5.23
<i>Piper sarmentosum</i> Roxb.	leaves	81.43 \pm 14.12
<i>Platycladus orientalis</i> (L.) Franco	leaves	92.41 \pm 16.49
<i>Plectranthus amboinicus</i> (Lour.) Spreng.	leaves	82.06 \pm 1.50
<i>Pluchea indica</i> (L.) Less.	leaves	88.07 \pm 4.37
<i>Pogostemon cablin</i> (Blanco) Benth.	leaves	93.64 \pm 1.51
<i>Vitex trifolia</i> L.	leaves	88.70 \pm 3.34
<i>Zingiber montanum</i> (J.Koenig) Link ex A.Dietr.	rhizomes	88.36 \pm 1.59
<i>Zingiber officinale</i> Roscoe	rhizomes	94.29 \pm 3.83
<i>Zingiber zerumbet</i> (L.) Roscoe ex Sm.	rhizomes	0.19 \pm 0.05

These selected EOs were further analyzed for dose-response activity to calculate IC₅₀ value and for cytotoxicity on mammalian cells. The results are summarized in Table 2. All selected EOs revealed high potential against promastigote form of *Lmm* with IC₅₀ values lower than 10 nL/mL. However, four of them also showed toxicity on mammalian cells, as indicated by their SI < 10. Our data demonstrated that the most active and selective EO was extracted from *Ocimum gratissimum* with an IC₅₀ value on *Lmm* promastigotes of 4.85 \pm 1.65 nL/mL and no toxicity towards mammalian cells at the highest tested concentration (50 nL/mL), as evidenced by more than 80% viable cells after 72 h of incubation.

Table 2. In vitro anti-leishmanial activity, cytotoxicity and selectivity indices of the five selected EOs.

Plant Species	Anti-Leishmanial Activity (IC ₅₀ nL/mL) Average \pm Standard Deviation	Cytotoxicity (IC ₅₀ nL/mL) Average \pm Standard Deviation			
		WI38	SI (WI38)	J774	SI (J774)
<i>Amomum aromaticum</i>	9.25 \pm 0.64	47.31 \pm 0.30	5.11	22.68 \pm 3.22	2.45
<i>Cinnamomum cassia</i>	2.92 \pm 0.08	14.19 \pm 0.54	4.85	6.26 \pm 0.80	2.14
<i>Elsholtzia ciliata</i>	8.49 \pm 0.32	47.38 \pm 1.64	5.58	13.21 \pm 1.48	1.56
<i>Ocimum gratissimum</i>	4.85 \pm 1.65	>50	>10.3	>50	>10.3
<i>Zingiber zerumbet</i>	3.34 \pm 0.34	3.68 \pm 0.34	1.10	2.41 \pm 0.04	0.72
Pentamidine	0.04 \pm 0.006 *				
Camptothecin		0.13 \pm 0.02 *		0.01 \pm 0.01 *	

* concentration in μ g/mL.

To define their composition and further determine active compounds, the five selected EOs were analyzed by GC-MS and GC-FID (Table 3). Identified compounds accounted for more than 90% for each EO. Eucalyptol (55.2%), *trans*-cinnamaldehyde (83.6%), citral (neral and geranial) (40.2%), eugenol (86.5%), and zerumbone (60.3%) were characterized as the main components of EOs extracted from *A. aromaticum*, *C. cassia*, *E. ciliata*, *O. gratissimum*, and *Z. zerumbet*, respectively.

Table 3. Chemical composition of the five selected EOs.

No.	Compounds	RI	Relative Percentages (%)					Identification
			<i>A. aromaticum</i>	<i>C. cassia</i>	<i>E. ciliata</i>	<i>O. gratissimum</i>	<i>Z. zerumbet</i>	
1	α -Pinene	535	2.1	-	-	-	2.3	MS, [10], Co-GC
2	Camphene	573–576	t	-	-	-	8.0	MS, [10]
3	β -Pinene	619–621	2.6	-	0.1	-	0.2	MS, [10], Co-GC
4	Sabinene	634–635	t	-	t	0.2	-	MS, [10]
5	3-Carene	664–665	0.1	-	-	-	1.1	MS
6	α -Phellandrene	682	-	-	-	-	0.1	MS, Co-GC
7	Myrcene	684	-	-	0.2	0.2	0.4	MS, [10]
8	α -Terpinene	697	-	-	-	t	-	MS, [10], Co-GC
9	Limonene	719–720	0.8	-	4.1	-	1.0	MS, [10], Co-GC
10	Eucalyptol	730-740	55.2	-	-	-	1.6	MS, [10], Co-GC
11	(Z)- β -Ocimene	758–760	-	-	0.7	5.4	-	MS
12	γ -Terpinene	765–766	-	-	t	0.1	t	MS, [10], Co-GC
13	(E)- β -Ocimene	773–778	-	-	14.0	0.2	-	MS
14	<i>p</i> -Cymene	787–792	0.6	t	t	-	0.2	MS, [10], Co-GC
15	Terpinolene	800–801	-	-	t	t	0.1	MS, [10], Co-GC
16	Octanal	812	0.2	-	-	-	-	MS
17	(E)-3-Hexen-1-ol acetate	837	-	-	t	-	-	MS
18	5-Hepten-2-one, 6-methyl-	855–858	0.2	-	1.1	-	-	MS, Co-GC
19	α -Pinene oxide	875	-	-	0.1	-	-	MS
20	<i>allo</i> -Ocimene	887	-	-	-	0.1	-	MS, Co-GC
21	1-Octen-1-yl acetate	897	-	-	1.0	-	-	MS
22	(Z)-3-Hexen-1-ol	902–904	-	-	0.5	t	-	MS, Co-GC
23	Fenchone	907	-	-	-	-	0.1	MS, Co-GC
24	3-Octanol	915	-	-	0.3	-	-	MS, [10]
25	(E)-2-Octenal	945	0.6	-	-	-	-	MS
26	<i>cis</i> -Linalool oxide (furanoid)	961	t	-	-	-	-	MS, Co-GC
27	1-Octen-3-ol	971	-	-	7.1	-	-	MS, [10]
28	<i>cis</i> -Sabinene hydrate	980	-	-	-	0.3	-	MS
29	Cyclosativene	987	-	0.1	-	-	-	MS
30	<i>trans</i> -Linalool oxide (furanoid)	988	t	-	-	-	-	MS, Co-GC
31	Citronellal	990	-	-	t	-	-	MS, [10], Co-GC
32	α -Copaene	998–1003	-	4.2	-	0.2	-	MS
33	Camphor	1020–1021	-	-	0.4	-	2.1	MS, [10], Co-GC
34	β -Bourbonene	1023	-	-	-	0.2	-	MS
35	Benzaldehyde	1027	-	1.0	t	-	-	MS
36	β -Cubebene	1044	-	-	-	0.1	-	MS
37	Linalool	1060–1065	0.7	-	8.3	0.1	0.3	MS, [10], Co-GC
38	Terpinen-1-ol	1077	0.1	-	-	-	-	MS
39	Bornyl acetate	1086	-	-	-	-	0.1	MS, Co-GC
40	<i>trans</i> - α -Bergamotene	1091–1092	-	t	t	-	-	MS, [10]
41	β -Elemene	1093–1094	-	0.2	-	0.1	-	MS
42	β -Caryophyllene	1098–1101	-	0.1	3.0	1.3	0.7	MS, [10], Co-GC
43	Terpinen-4-ol	1108–1113	0.9	t	t	0.2	0.2	MS, [10], Co-GC
44	Acetophenone	1149	-	-	0.7	-	-	MS
45	(E)-2-Decenal	1153	5.3	-	-	-	-	MS

Table 3. Cont.

No.	Compounds	RI	Relative Percentages (%)					Identification
			<i>A. aromaticum</i>	<i>C. cassia</i>	<i>E. ciliata</i>	<i>O. gratissimum</i>	<i>Z. zerumbet</i>	
46	α -Humulene	1166–1173	-	t	t	t	9.2	MS, [10], Co-GC
47	(<i>E</i>)- β -Farnesene	1178	-	-	6.2	-	-	MS
48	γ -Muuroleone	1187–1188	-	0.5	-	t	-	MS
49	Neral	1187–1188	3.2	-	16.8	-	-	MS, Co-GC
50	Methyl geranate	1197	-	-	0.2	-	-	MS
51	α -Terpineol	1201–1207	2.8	-	0.4	-	0.2	MS, [10], Co-GC
52	Borneol	1205	-	-	-	-	0.2	MS, [10], Co-GC
53	Germacrene D	1207	-	-	-	3.4	-	MS, [10]
54	Geranyl formate	1212	0.6	-	-	-	-	MS
55	β -Selinene	1214	-	0.1	-	-	-	MS
56	Neryl acetate	1218	-	-	t	-	-	MS, [10], Co-GC
57	α -Muuroleone	1224	-	0.8	-	-	-	MS
58	Geranial	1238	6.4	-	23.4	-	-	MS
59	β -Cadinene	1253–1254	-	-	-	0.2	0.1	MS
60	δ -Cadinene	1257	-	2.0	-	-	-	MS
61	Geranyl acetate	1259–1263	0.9	-	0.7	-	-	MS, [10], Co-GC
62	Citronellol	1272	-	-	0.5	-	-	MS, [10], Co-GC
63	Benzenepropanal	1273	-	1.7	-	-	-	MS
64	<i>cis</i> -Cadin-1,4-diene	1278	-	0.5	-	-	-	MS
65	Isogeraniol	1289	-	-	0.2	-	-	MS
66	Nerol	1304	-	-	3.5	-	-	MS, [10], Co-GC
67	<i>cis</i> -Isogeraniol	1311	-	-	0.4	-	-	MS
68	(<i>E</i>)-2,6-dimethyl-3,5,7-octatriene-2-ol	1323–1324	-	-	t	0.1	-	MS
69	<i>cis</i> -Calamenene	1325	-	0.5	-	-	-	MS
70	Geraniol	1350–1354	2.4	-	3.1	-	-	MS, [10], Co-GC
71	(<i>E</i>)-2-Dodecenal	1359	1.5	-	-	-	-	MS
72	<i>cis</i> -Cinnamaldehyde	1382	-	2.4	-	-	-	MS
73	Indane-4-carboxaldehyde	1466	1.7	-	-	-	-	MS
74	Caryophyllene oxide	1466–1501	-	-	0.6	0.2	3.6	MS, [10], Co-GC
75	Humulene epoxide II	1522	-	-	-	-	2.2	MS
76	<i>trans</i>-Cinnamaldehyde	1534	-	83.6	-	-	-	MS, Co-GC
77	(<i>E</i>)-Nerolidol	1533–1538	1.2	-	0.5	-	-	MS, [10], Co-GC
78	Cubenol	1552	-	0.2	-	-	-	MS
79	Elemol	1571	0.2	-	-	-	-	MS
80	Cinnamyl acetate	1633	-	0.9	-	-	-	MS
81	Eugenol	1657	-	-	-	86.5	-	MS, Co-GC
82	α -Muurolol	1669	-	0.1	-	-	-	MS
83	β -Eudesmol	1706	-	-	-	-	0.2	MS
84	α -Cadinol	1710	-	-	-	0.1	-	MS
85	Cinnamyl alcohol	1761	-	0.2	-	-	-	MS, Co-GC
86	Zerumbone	1831	-	-	-	-	60.3	MS
87	<i>o</i> -Methoxy-cinnamaldehyde	1902	-	0.3	-	-	-	MS
88	<i>trans</i> -Phytol	2085	-	-	0.16	-	-	MS, Co-GC
	Total identified		90.3	99.4	98.3	99.2	94.5	

t: trace (peak area less than 0.05%); RI: the retention index was calculated using a homologous series of fatty acid methyl esters C₅–C₂₇; MS: mass spectra (matching coefficient >700 compared with NIST database); Co-GC: co-injection with pure compound.

It is important to point out that eugenol is the major compound (86.5%) of the EO extracted from *O. gratissimum*, which was shown to be the most selective active sample. In order to understand the activity of this EO, in vitro anti-leishmanial activity and cytotoxicity of eugenol were studied. Interestingly, the IC₅₀ value against *Lmm* promastigotes of eugenol was 2.57 ± 0.57 nL/mL ($=2.72$ µg/mL, 15.67 µM) and more than 80% of mammalian cells were living after 72 h when incubated at the maximal concentration used (50 nL/mL).

3. Discussion

This study analyzed the in vitro activity of 37 EOs extracted from Vietnamese plants against *Leishmania mexicana mexicana*. According to the classification of anti-parasitic activity from literature, extracts that showed effect against parasites with IC₅₀ value ≤ 2 µg/mL (or 2 µM for pure compounds) are cited as having good activity, those with IC₅₀ between 2 and 20 µg/mL (or micro molar for pure compounds) are considered as having a moderate activity, while those with higher IC₅₀ value were considered as less interesting [11]. In this study, the primary screening was performed against *Lmm* promastigotes at concentrations of 25 and 50 nL/mL in order to quickly identify interesting EOs (1 nL is a little less than 1 µg depending on the density of the EO) [11]. Our data showed that percentages of viable parasites incubated with the lower concentration of five EOs, extracted from *A. aromaticum*, *C. cassia*, *E. ciliata*, *O. gratissimum*, and *Z. zerumbet*, were lower than 1%, meaning that these EOs revealed a potential activity. Indeed, the IC₅₀ values of these selected EOs were lower than 10 nL/mL in the second analysis.

To have a general look at anti-leishmanial activity and cytotoxicity of the five more active EOs and to understand their observed effects, we compiled literature data of EOs extracted from similar plant species and also of their major components in Table 4.

Table 4. Profile of in vitro anti-leishmanial activity and cytotoxicity of the five more active EOs and their major compounds.

Plants/Compounds	Anti-Leishmanial Activity		Cytotoxicity		Refs.	
	L. Species-Form	IC ₅₀ (µg/mL)	Cell Line	IC ₅₀ (µg/mL)		
<i>A. aromaticum</i>	ND		ND			
<i>C. cassia</i>	ND		ND			
<i>E. ciliata</i>	ND		PC12 rat pheochromocytoma cells	>50	[12]	
<i>O. gratissimum</i>	<i>L. amazonensis</i> -proma.	135	CHO	125.00 ± 1.68	[13,14]	
	<i>L. amazonensis</i> -ama.	100	WI38	165.51 ± 6.81	[13,14]	
	<i>L. chagasi</i> -proma.	80			[15]	
<i>Z. zerumbet</i>	<i>L. donovani</i> -proma.	4.62	ND		[16]	
Eucalyptol	<i>L. infantum</i> -proma.	>100	Vero cell	63.49	[17]	
	<i>L. infantum</i> -ama.	>100			[17]	
	<i>L. infantum</i> -proma.	>400			[18]	
	<i>L. tropica</i> -proma.	>400			[18]	
	<i>L. major</i> -proma.	>400			[18]	
Cinnamal-dehyde	ND		human embryonic stem cells	4.88	[19]	
			human pulmonary fibroblasts	5.28	[19]	
Citral	<i>L. donovani</i> -proma.	19	kidney epithelial cell J774	22.4	[20]	
	<i>L. amazonensis</i> -proma.	8.0 ± 0.06			50.0 ± 0.10	[21]
	<i>L. amazonensis</i> -ama.	25 ± 0.29				[21]
	<i>L. infantum</i> -proma.	42				[22]
	<i>L. tropica</i> -proma.	34				[22]
	<i>L. major</i> -proma.	36				[22]
Eugenol	<i>L. amazonensis</i> -proma.	12.65	red blood cell BALB/c peritoneal macrophages	>65.6		[23]
	<i>L. infantum chagasi</i> -proma.	500			300	[24]
	<i>L. infantum chagasi</i> -axe. ama.	220				[24]
	<i>L. infantum chagasi</i> infected macrophages	100				[24]
	<i>L. infantum chagasi</i> -proma.	56.13 ± 2.09				[25]
	<i>L. infantum chagasi</i> -ama.	20.81 ± 1.59				[25]

Table 4. Cont.

Plants/Compounds	Anti-Leishmanial Activity		Cytotoxicity		Refs.
	L. Species-Form	IC ₅₀ (µg/mL)	Cell Line	IC ₅₀ (µg/mL)	
Eugenol (emulsified)	<i>L. donovani</i> -proma.	8.43 ± 0.96	murine macrophages	>200	[26]
	<i>L. donovani</i> -ama.	5.05 ± 1.72			[26]
Zerumbone	<i>L. donovani</i> -proma.	2.04	HL-60	2.27	[27,28]

ND: not determined; proma.: promastigote; ama.: intracellular amastigote; axe. ama.: axenic amastigote.

As shown in this table, among the five selected samples, EOs obtained from *A. aromaticum*, *C. cassia*, and *E. ciliata* were analyzed here for the first time for their leishmanicidal effect. In our experiment, they showed IC₅₀ values of 9.25 ± 0.64, 2.29 ± 0.08, and 8.49 ± 0.32 nL/mL, respectively against *Lmm* promastigotes. Unfortunately, this activity was not very selective on *Leishmania* as SI values compared to non-cancer mammalian cells (WI38) and cancer cells (J774) were approximately 5 and 2, respectively.

Two EOs extracted from *O. gratissimum* and *Z. zerumbet* were already investigated for anti-leishmanial activity [13–16] however on other *L.* species. The EO extracted from fresh leaves of *O. gratissimum* collected in Brazil did not reveal interesting effects against both *L. amazonensis* and *L. chagasi*. On the contrary, in our study, the EO extracted from *O. gratissimum* collected in Vietnam demonstrated notable activity against *Lmm* promastigotes, with an IC₅₀ value of 4.85 ± 1.65 nL/mL. These results indicated a specific activity of this EO related to parasite species (*Leishmania mexicana mexicana*) and/or collection place and composition. Another important feature of this EO is its absence of cytotoxicity as the percentages of viable WI38 and J774 cells at the maximal tested concentration (50 nL/mL) were higher than 80%. We therefore selected the EO extracted from *O. gratissimum* as the most active and selective sample. Regarding the EO extracted from fresh rhizomes of *Z. zerumbet*, its activity against *L. donovani* was in the same range than what we found against *Lmm* promastigotes (IC₅₀ = 3.34 ± 0.34 nL/mL). Unfortunately, we also observed cytotoxicity of this EO on WI38 and J774 cells as shown by SI values of 1.10 and 0.72, respectively.

Eucalyptol, the major compound of the EO extracted from *A. aromaticum* (55.2%), was not active against the different tested *L.* species [17,18]. Other components present in our *A. aromaticum* EO, such as citral (neral and geranial, 9.6%), 2-(*E*)-decenal (5.3%), α-terpineol (2.8%), β-pinene (2.6%) and α-pinene (2.1%) may be therefore responsible, in part, of the observed activity. Among these compounds, 2-(*E*)-decenal and α-pinene have already shown leishmanicidal effects. IC₅₀ values of 2-(*E*)-decenal against *L. donovani* promastigotes and axenic amastigotes were 7.85 ± 0.28 and 2.47 ± 0.25 µg/mL respectively [29], and IC₅₀ value of α-pinene against *L. amazonensis* promastigotes, *L. amazonensis* axenic amastigotes and *L. infantum* promastigotes were 19.7, 16.1 and 45.94 µg/mL respectively [17,30].

Citral was characterized as the main component of the EO extracted from *E. ciliata* (40.2%). It showed a moderate activity against *L. donovani*, *L. amazonensis* and a weak effect against *L. infantum*, *L. tropica* and *L. major* [20–22]. However, it is important to point out that this compound is not selective on *Leishmania* as shown by the toxicity on kidney epithelial and J774 cells [20,21]. Our results indicated activity on parasites and mammalian cells of the EO extracted from *E. ciliata* in the same range than citral with IC₅₀ value of 8.49 ± 0.32 nL/mL and SI of 5.58 (compared to WI38) and 1.56 (compared to J774). Nevertheless, citral being present at only 40.2% in the *E. ciliata* EO, other compounds should also have anti-leishmanial activity as β-(*E*)-ocimene (14.0%), linalool (8.3%), 1-octen-3-ol (7.1%) and β-(*E*)-farnesene (6.2%). Dutra et al. have already examined leishmanicidal effect of linalool on *L. infantum chagasi*. However when axenic amastigotes were treated with linalool, IC₅₀ was 550 µg/mL [24].

EO extracted from *C. cassia* contained 83.6% of *trans*-cinnamaldehyde. The anti-leishmanial activity of this compound is unknown but its toxicity on two human cell lines was determined [19]. In our sample, the high percentage of cinnamaldehyde may explain the toxicity of this EO on both mammalian cell lines, WI38 and J774, with IC₅₀ of 14.19 ± 0.54 and 6.26 ± 0.80 nL/mL respectively.

Given the known cytotoxicity of this compound and its high concentration in the *C. cassia* EO, we did not find interesting to further analyze its anti-parasitic activity.

As mentioned previously, eugenol was characterized as the major compound (86.5%) of the most interesting EO extracted from our Vietnamese sample of *O. gratissimum*. Literature data indicated a moderate effect on *L. amazonensis* and a less interesting activity against *L. infantum chagasi* of eugenol [23–25]. On *L. donovani*, along with a moderate in vitro activity, in vivo effect of eugenol (emulsified) was determined. The intra-peritoneal administration of an eugenol emulsion at the dose of 75 mg/kg b.w. for 10 consecutive days decreased by 87.01 ± 5.85 and $86.68 \pm 5.42\%$ parasitic load in spleen and liver, respectively, in 8-weeks infected BALB/c mice. Moreover, a significant reduction in spleen size, and spleen and liver weights were also found at this dose [26]. Our results further support the anti-leishmanial activity of this compound with IC_{50} value of 2.57 ± 0.57 nL/mL ($=2.72$ μ g/mL or 15.67 μ M) against *Lmm* promastigotes. Interestingly, eugenol was not toxic on both non-cancer and cancer mammalian cells in our models at the highest analyzed concentration (50 nL/mL). From these results, and its high percentage in the *O. gratissimum* EO, it can be concluded that activity of eugenol can explain the anti-leishmanial activity of this EO.

Zerumbone, accounting for 60.3% of the EO extracted from fresh rhizomes of *Z. zerumbet*, was reported to be active against *L. donovani*, however it revealed toxicity on human leukemia cells HL-60 [27,28]. These data can explain the anti-leishmanial activity but also cytotoxicity of the *Z. zerumbet* EO found in our study.

Mechanisms of anti-leishmanial activity of these EOs and their major compounds are not well known. Usually, effects were analyzed on the morphology of treated parasites or as a consequence of immunostimulatory activities.

Most of the studies used transmission electron microscopy (TEM) and scanning electron microscopy (SEM) to analyze the morphology of EOs treated parasites. The EO extracted from *O. gratissimum* caused considerable mitochondrial swelling in *L. amazonensis* promastigotes at 135 μ g/mL and amastigotes at 100 μ g/mL [13], swelling of cell body, flagellar pocket and mitochondria in *L. chagasi* promastigotes at 50 μ g/mL [15]. Ultrastructure alterations were also observed in *L. amazonensis* and *L. infantum* promastigotes treated with citral at concentrations of 8.0 and 42 μ g/mL respectively [21,22]. Mukherjee et al. detected morphological alterations by SEM in *L. donovani* promastigotes treated with 9.36 μ M of zerumbone associated with induced ROS-mediated apoptosis [27].

Using other experiments, citral at the concentration of 42 μ g/mL was shown to trigger programmed cell death of *L. infantum* promastigotes, as indicated by the externalization of phosphatidylserine, loss of mitochondrial membrane potential and cell-cycle arrest at the G(0)/G(1) phase [22].

Regarding immunostimulatory activity, an increase of nitric oxide produced by infected macrophages has been suggested to be responsible for the activity of the *O. gratissimum* EO against *L. amazonensis* at 100 and 150 μ g/mL [13]. Islamuddin et al. explored the synergic effect between eugenol emulsion and the immune system. In BALB/mice infected by *L. donovani*, treatment with 75 mg/kg b.w. of eugenol emulsion enhanced IFN- γ and IL-2 serum levels, as well as increased CD4+ and CD8+ T cell population and expanded IFN- γ producing CD4+ and CD8+ splenic T lymphocytes [26].

Concerning the mechanism of eugenol activity, although no data are available on *Lmm* promastigotes, some experiments were carried out on bacteria and fungi. Eugenol at concentrations of 5.3 and 10.6 mg/mL was reported to significantly damage both the cell wall and membrane of the treated Gram-negative and Gram-positive bacteria [31]. Khan et al. observed damaging effects of eugenol at 200 μ g/mL on cell wall, cell membrane, cytoplasmic contents and other membranous structures of treated *Candida albicans* [32]. Indeed, because of the lipophilicity of eugenol, it could easily diffuse between the fatty acyl chains of lipid bilayers modifying the fluidity, integrity and

permeability of cell membranes [31]. We now intend to analyze possible effects of this compound on leishmanial membranes.

4. Materials and Methods

4.1. Plants Collection

Thirty-seven plants were collected from different areas of Vietnam in November 2014 and from May to August 2015. They were identified by matching with literature and herbarium specimen at the Botanical Department, Hanoi University of Pharmacy, Vietnam. The information of sample name, genus, species, family, and collector name is given in the supplementary material.

4.2. Essential Oils Extraction

Fresh samples were extracted by hydro-distillation using a modified Clavenger apparatus for two hours. Each essential oil obtained from hydro-distillation was dried using sodium sulfate, filtered and kept refrigerated prior to analysis. Stock solutions at the concentration of 20 $\mu\text{L}/\text{mL}$ were prepared in DMSO and diluted further in culture medium to achieve a maximum final DMSO concentration of 0.25%.

4.3. Culture Maintenance

Promastigote form of *Leishmania mexicana mexicana* (*Lmm*, MHOM/BZ/84/BEL46) was grown in SDM 79 medium (Life technologies, Ghent, Belgium) supplemented with 15% heat-inactivated fetal bovine serum and 0.2% hemin. The culture was maintained at 28 °C in 5% CO₂ incubator. The human non-cancer fibroblast cell line WI38 (ATCC Number CCL-75 from LGC Standards, Middlesex, UK) and macrophage-like murine cell line J774 (ECACC Number 91051511 from Public Health England, Salisbury, UK) were grown in DMEM and RPMI medium (Gibco from Thermo Fisher Scientific, Merelbeke, Belgium or Sigma-Aldrich, Bornem, Belgium), respectively, supplemented with 10% fetal bovine serum and penicillin-streptomycin (100 UI/mL). The culture was maintained at 37 °C in 5% CO₂ incubator.

4.4. Anti-leishmanial Assay

The *Lmm* promastigote density was counted on a haemocytometer and adjusted to 10⁵ parasites/mL. The assays were performed in 96-well plates. For primary screening (repeated two times at concentrations of 50 and 25 nL/mL in triplicate) essential oils were diluted in culture medium from the stock solutions (20 $\mu\text{L}/\text{mL}$). Each well was filled with 50 μL of the diluted essential oil and 50 μL of the *Lmm* promastigote culture (total volume 100 μL). After 72 h of incubation, 10 μL Alamar blue (Thermo Fisher Scientific, diluted with PBS at the ratio 1:1) was added to each well and the plates were further incubated for 4 h. Fluorescence was measured on a spectrophotometer (SpectraMax-Molecular Devices, Berkshire, UK) at 530 nm excitation and 590 nm emission wavelengths. Pentamidine was tested as standard drug. The essential oils that inhibited more than 50% the growth of *Lmm* promastigotes at the concentration of 25 nL/mL in the primary screening were submitted to a second analysis for accurate IC₅₀ determination. Selected essential oils were screened at least three times at concentrations ranging from 50 to 0.02 nL/mL in duplicate. IC₅₀ values were calculated from dose response growth inhibition curves by Microsoft excel files.

4.5. Cytotoxicity Assay

The essential oils which were analyzed for IC₅₀ value on anti-leishmanial assay were also analyzed for cytotoxicity against non-cancer (WI38) and cancer (J774) mammalian cells. The culture of cells was diluted with medium to the adequate density of 5 × 10³ cells/mL and then 180 μL was added into each well of 96-well plates. After 24 h of incubation, 20 μL of essential oils diluted in culture medium was added to each well (to obtain concentrations ranging from 50–0.02 nL/mL) for further 72 h of

incubation. After removing the medium, 100 μ L of MTT was added to each well and the plates were further incubated for 45 minutes. 100 μ L of DMSO was used to dissolve formed formazan crystals after MTT removing. Absorbance was measured on a spectrophotometer (SpectraMax-Molecular Devices, Berkshire, UK) at 570 nm with a reference wavelength at 620 nm. All experiments were made at least two times in triplicate. The computation of the IC₅₀ values was performed with GraphPad Prism 5.0 (GraphPad Software Inc., San Diego, CA, USA). The selectivity index (SI) values were calculated using the formula below:

$$SI = IC_{50} \text{ for mammalian cell} / IC_{50} \text{ for protozoan parasite}$$

4.6. Essential Oil Analysis

The GC-MS analyses were carried out on a TRACE GC 2000 series (Thermo-Quest, Rodano, Italy), with a DB-WAX capillary column (30 m \times 0.25 mm \times 0.25 μ m) using the following operating conditions: injection volume: 1 μ L (TBME solution); injection mode: splitless; injector temperature: 230 $^{\circ}$ C; oven temperature: increased from 45 $^{\circ}$ C (held on 5 min) to 250 $^{\circ}$ C (held on 5 min) at 3 $^{\circ}$ C/min; helium was used as a carrier gas at a constant flow of 1.3 mL/min; detector temperatures: 260 $^{\circ}$ C; ion source: 70 eV. The oil components were identified using linear retention indices in relation to a series of fatty acid methyl esters (C₅–C₂₇), pure compounds and NIST mass spectral library (matching coefficient > 700). GC-FID was done using the same column and conditions on a FOCUS GC (Thermo Finnigan, Milan, Italy) with modifications of injection mode (split at ratio 1:50) and detector temperature (250 $^{\circ}$ C). Percentage of compounds was calculated by the normalization procedure.

5. Conclusions

The present study analyzes for the first time the anti-leishmanial activity of 37 essential oils extracted from Vietnamese plants. Those extracted from *A. aromaticum*, *C. cassia* and *E. ciliata* are shown here for the first time to be effective on a *Leishmania* species, while the effects of *O. gratissimum* and *Z. zerumbet* EOs are reported for the first time on *L. mexicana* species. More than 90% of their contents was characterized. Eugenol was identified as the major compound of the *O. gratissimum* EO, the most active and selective one. This compound, showing a moderate anti-leishmanial activity and low cytotoxicity in the tested models, can explain this EO activity. However, further results are necessary before developing it in the treatment of cutaneous leishmaniasis such as in vivo assessment in *Lmm* infected animals and determination of its mode of action.

Supplementary Materials: Supplementary materials are available online.

Acknowledgments: The authors gratefully acknowledge Vietnamese Ministry of Education and Training and Belgian National Fund for Scientific Research (FNRS) (PDR T.0190.13) for financial support. We wish to thank Marie-France Hérent and Marie-Christine Fayt for skillful technical assistance in GC and all the members of the Pharmacognosy group from LDRI, UCL, Belgium, for helpful scientific advices and support.

Author Contributions: Thanh Binh Le directly collected plants, performed chemical, anti-leishmanial activity and cytotoxicity experiments, analyzed the data; Claire Beaufay participated in analyzes of the data and discussions during the work; Duc Trong Nghiem was responsible for the plants identification; Marie-Paule Mingeot-Leclercq participated in discussions during the work; Joëlle Quetin-Leclercq was responsible for the conception of this work. They also wrote this article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Centers for Disease Control and Prevention (CDC). Parasites-Leishmaniasis. Available online: <https://www.cdc.gov/parasites/leishmaniasis/> (accessed on 8 June 2017).
2. WHO Expert Committee. *Control of the Leishmaniases*; WHO Technical Report Series, No. 949; World Health Organization: Geneva, Switzerland, 2010.

3. Drugs for Neglected Diseases initiative (DNDi). Diseases & Projects-Leishmaniasis. Available online: <http://www.dndi.org/diseases-projects/leishmaniasis/> (accessed on 8 June 2017).
4. Schmidt, T.J.; Khalid, S.A.; Romanha, A.J.; Alves, T.M.; Biavatti, M.W.; Brun, R.; da Costa, F.B.; de Castro, S.L.; Ferreira, V.F.; de Lacerda, M.V.; et al. The potential of secondary metabolites from plants as drugs or leads against protozoan neglected diseases—part I. *Curr. Med. Chem.* **2012**, *19*, 2128–2175. [[CrossRef](#)] [[PubMed](#)]
5. Bakkali, F.; Averbeck, S.; Averbeck, D.; Idaomar, M. Biological effects of essential oils—A review. *Food. Chem. Toxicol.* **2008**, *46*, 446–475. [[CrossRef](#)] [[PubMed](#)]
6. Monzote, L.; García, M.; Pastor, J.; Gil, L.; Scull, R.; Maes, L.; Cos, P.; Gille, L. Essential oil from *Chenopodium ambrosioides* and main components: Activity against *Leishmania*, their mitochondria and other microorganisms. *Exp. Parasitol.* **2014**, *136*, 20–26. [[CrossRef](#)] [[PubMed](#)]
7. Monzote, L.; Pastor, J.; Scull, R.; Gille, L. Antileishmanial activity of essential oil from *Chenopodium ambrosioides* and its main components against experimental cutaneous leishmaniasis in BALB/c mice. *Phytomedicine* **2014**, *21*, 1048–1052. [[CrossRef](#)] [[PubMed](#)]
8. Monzote, L.; García, M.; Scull, R.; Cuellar, A.; Setzer, W.N. Antileishmanial activity of the essential oil from *Bixa orellana*. *Phytother. Res.* **2014**, *28*, 753–758. [[CrossRef](#)] [[PubMed](#)]
9. Montoro, P.; Masullo, M.; Piacente, S.; Pizza, C. Extraction, sample preparation, and analytical methods for quality issues of essential oils. In *Aromatherapy: Basic Mechanisms and Evidence-Based Clinical Use*; Bagetta, G., Cosentino, M., Sakurada, T., Eds.; CRC Press: Boca Raton, FL, USA, 2015; p. 106, ISBN 978-1-4822-4663-6.
10. Hérent, M.F.; de Bie, V.; Tilquin, B. Determination of new retention indices for quick identification of essential oils compounds. *J. Pharm. Biomed. Anal.* **2007**, *43*, 886–892. [[CrossRef](#)] [[PubMed](#)]
11. Bero, J.; Kpoviessi, S.; Quetin-Leclercq, J. Anti-parasitic activity of essential oils and their constituents against *Plasmodium*, *Trypanosoma* and *Leishmania*. In *Novel Plant Bioresource: Applications in Food, Medicine and Cosmetic*; Gurib-Fakim, A., Ed.; John Wiley & Sons: Oxford, UK, 2014; pp. 445–469, ISBN 978-1-118-46061-0.
12. Choi, M.S.; Choi, B.S.; Kim, S.H.; Pak, S.C.; Jang, C.H.; Chin, Y.W.; Kim, Y.M.; Kim, D.I.; Jeon, S.; Koo, B.S. Essential Oils from the Medicinal Herbs Upregulate Dopamine Transporter in Rat Pheochromocytoma Cells. *J. Med. Food.* **2015**, *18*, 1112–1120. [[CrossRef](#)] [[PubMed](#)]
13. Ueda-Nakamura, T.; Mendonça-Filho, R.R.; Morgado-Díaz, J.A.; Korehisa Maza, P.; Prado Dias Filho, B.; Aparício Garcia Cortez, D.; Alviano, D.S.; Rosa Mdo, S.; Lopes, A.H.; Alviano, C.S.; et al. Antileishmanial activity of Eugenol-rich essential oil from *Ocimum gratissimum*. *Parasitol. Int.* **2006**, *55*, 99–105. [[CrossRef](#)] [[PubMed](#)]
14. Kpadonou Kpoviessi, B.G.H.; Kpoviessi, D.S.S.; Yayi Ladekan, E.; Gbaguidi, F.; Frédérick, M.; Moudachirou, M.; Quetin-Leclercq, J.; Accrombessi, G.C.; Bero, J. In vitro antitrypanosomal and antiplasmodial activities of crude extracts and essential oils of *Ocimum gratissimum* Linn from Benin and influence of vegetative stage. *J. Ethnopharmacol.* **2014**, *155*, 1417–1423. [[CrossRef](#)] [[PubMed](#)]
15. Oliveira, V.C.S.; Moura, D.M.S.; Lopes, J.A.D.; de Andrade, P.P.; da Silva, N.H.; Figueiredo, R.C.B.Q. Effects of essential oils from *Cymbopogon citratus* (DC) Stapf., *Lippia sidoides* Cham., and *Ocimum gratissimum* L. on growth and ultrastructure of *Leishmania chagasi* promastigotes. *Parasitol. Res.* **2009**, *104*, 1053–1059. [[CrossRef](#)] [[PubMed](#)]
16. Singh, C.B.; Chanu, S.B.; Kh, L.; Swapana, N.; Cantrell, C.; Ross, S.A. Chemical composition and biological activity of the essential oil of rhizome of *Zingiber zerumbet* (L.) Smith. *J. Pharmacogn. Phytochem.* **2014**, *3*, 130–133. Available online: <http://handle.nal.usda.gov/10113/62328> (accessed on 8 June 2017).
17. Leal, S.M.; Pino, N.; Stashenko, E.E.; Martínez, J.R.; Escobar, P. Antiprotozoal activity of essential oils derived from *Piper* spp. grown in Colombia. *J. Essent. Oil Res.* **2013**, *25*, 512–519. [[CrossRef](#)]
18. Machado, M.; Dinis, A.M.; Santos-Rosa, M.; Alves, V.; Salgueiro, L.; Cavaleiro, C.; Sousa, M.C. Activity of *Thymus capitellatus* volatile extract, 1,8-cineole and borneol against *Leishmania* species. *Vet. Parasitol.* **2014**, *200*, 39–49. [[CrossRef](#)] [[PubMed](#)]
19. Behar, R.Z.; Davis, B.; Wang, Y.; Bahl, V.; Lin, S.; Talbot, P. Identification of toxicants in cinnamon-flavored electronic cigarette refill fluids. *Toxicol. In Vitro* **2014**, *28*, 198–208. [[CrossRef](#)] [[PubMed](#)]
20. Zheljzkov, V.D.; Cantrell, C.L.; Tekwani, B.; Khan, S.I. Content, composition, and bioactivity of the essential oils of three basil genotypes as a function of harvesting. *J. Agric. Food Chem.* **2008**, *56*, 380–385. [[CrossRef](#)] [[PubMed](#)]

21. Santin, M.R.; dos Santos, A.O.; Nakamura, C.V.; Dias Filho, B.P.; Ferreira, I.C.P.; Ueda-Nakamura, T. In vitro activity of the essential oil of *Cymbopogon citratus* and its major component (citral) on *Leishmania amazonensis*. *Parasitol. Res.* **2009**, *105*, 1489–1496. [[CrossRef](#)] [[PubMed](#)]
22. Machado, M.; Pires, P.; Dinis, A.M.; Santos-Rosa, M.; Alves, V.; Salgueiro, L.; Cavaleiro, C.; Sousa, M.C. Monoterpenic aldehydes as potential anti-*Leishmania* agents: Activity of *Cymbopogon citratus* and citral on *L. infantum*, *L. tropica* and *L. major*. *Exp. Parasitol.* **2012**, *130*, 223–231. [[CrossRef](#)] [[PubMed](#)]
23. Xavier, F.J.S.; Rodrigues, K.A.D.F.; de Oliveira, R.G.; Lima Junior, C.G.; Rocha, J.D.; Keesen, T.S.; de Oliveira, M.R.; Silva, F.P.; Vasconcelos, M.L. Synthesis and in vitro anti *Leishmania amazonensis* biological screening of Morita-Baylis-Hillman Adducts prepared from eugenol, thymol and carvacrol. *Molecules* **2016**, *21*, 1483. [[CrossRef](#)] [[PubMed](#)]
24. Dutra, F.L.; Oliveira, M.M.; Santos, R.S.; Silva, W.S.; Alviano, D.S.; Vieira, D.P.; Lopes, A.H. Effects of linalool and eugenol on the survival of *Leishmania (L.) infantum chagasi* within macrophages. *Acta Trop.* **2016**, *164*, 69–76. [[CrossRef](#)] [[PubMed](#)]
25. De Moraes, S.M.; Vila-Nova, N.S.; Bevilaqua, C.M.L.; Rondon, F.C.; Lobo, C.H.; de Alencar Araripe Noronha Moura, A.; Sales, A.D.; Rodrigues, A.P.; de Figueiredo, J.R.; Campello, C.C.; et al. Thymol and eugenol derivatives as potential antileishmanial agents. *Bioorg. Med. Chem.* **2014**, *22*, 6250–6255. [[CrossRef](#)] [[PubMed](#)]
26. Islamuddin, M.; Chouhan, G.; Want, M.Y.; Ozbak, H.A.; Hemeg, H.A.; Afrin, F. Immunotherapeutic potential of eugenol emulsion in experimental visceral leishmaniasis. *PLoS Negl. Trop. Dis.* **2016**, *10*, e0005011. [[CrossRef](#)] [[PubMed](#)]
27. Mukherjee, D.; Singh, C.B.; Dey, S.; Mandal, S.; Ghosh, J.; Mallick, S.; Hussain, A.; Swapana, N.; Ross, S.A.; Pal, C. Induction of apoptosis by zerumbone isolated from *Zingiber zerumbet* (L.) Smith in protozoan parasite *Leishmania donovani* due to oxidative stress. *Braz. J. Infect. Dis.* **2016**, *20*, 48–55. [[CrossRef](#)] [[PubMed](#)]
28. Huang, N.C.; Chien, T.Y.; Chen, L.G.; Wang, C.C. Antitumor effects of zerumbone from *Zingiber zerumbet* in P-388D1 cells in vitro and in vivo. *Planta Med.* **2005**, *71*, 219–224. [[CrossRef](#)] [[PubMed](#)]
29. Donega, M.A.; Mello, S.C.; Moraes, R.M.; Jain, S.K.; Tekwani, B.L.; Cantrell, C.L. Pharmacological activities of cilantro's aliphatic aldehydes against *Leishmania donovani*. *Planta Med.* **2014**, *80*, 1706–1711. [[CrossRef](#)] [[PubMed](#)]
30. Rodrigues, K.A.D.F.; Amorim, L.V.; Dias, C.N.; Moraes, D.F.C.; Carneiro, S.M.P.; Carvalho, F.A.D.A. *Syzygium cumini* (L.) Skeels essential oil and its major constituent α -pinene exhibit anti-*Leishmania* activity through immunomodulation in vitro. *J. Ethnopharmacol.* **2015**, *160*, 32–40. [[CrossRef](#)] [[PubMed](#)]
31. Kamatou, G.P.; Vermaak, I.; Viljoen, A.M. Eugenol-From the remote Maluku Islands to the international market place: A review of a remarkable and versatile molecule. *Molecules* **2012**, *17*, 6953–6981. [[CrossRef](#)] [[PubMed](#)]
32. Khan, M.S.A.; Ahmad, I.; Cameotra, S.S. Phenyl aldehyde and propanoids exert multiple sites of action towards cell membrane and cell wall targeting ergosterol in *Candida albicans*. *AMB Express* **2013**, *3*, 54. [[CrossRef](#)] [[PubMed](#)]

Sample Availability: Samples of essential oils are available from the authors.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).



HANOI UNIVERSITY OF PHARMACY, VIETNAM

DEPARTMENT OF BOTANY

HERBARIUM OF MEDICINAL PLANTS (HNIP)

CERTIFICATE OF NUMBER SIGN OF SAMPLES

No.	Sign	Collector & Sender	Date of collection	Date sent	Scientific name	Vietnamese name	Locality	Determined by	Duplicate	Number sign in HNIP
1	TD_01	Le Thanh Binh	05/2015	08/08/2016	<i>Melaleuca alternifolia</i> (Maiden & Betch) Cheel	Tràm Úc	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18250/16
2	TD_02	Le Thanh Binh	05/2015	08/08/2016	<i>Ocimum gratissimum</i> L.	Hương nhu trắng	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18251/16
3	TD_03	Le Thanh Binh	05/2015	08/08/2016	<i>Pluchea indica</i> (L.) Less.	Cúc tần	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18252/16
4	TD_04	Le Thanh Binh	05/2015	08/08/2016	<i>Platyclusus orientalis</i> (L.) Franco	Trắc bách	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18253/16
5	TD_05	Le Thanh Binh	06/2015	08/08/2016	<i>Piper sarmentosum</i> Roxb.	Lá lốt	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18254/16
6	TD_06	Le Thanh Binh	08/2015	08/08/2016	<i>Artemisia annua</i> L.	Thanh hao hoa vàng	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18255/16
7	TD_07	Le Thanh Binh	05/2015	08/08/2016	<i>Ocimum tenuiflorum</i> L.	Hương nhu tía	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18256/16
8	TD_08	Le Thanh Binh	08/2015	08/08/2016	<i>Dysphania ambrosioides</i> (L.) Mosyakin & Clemant	Dầu giun	Thanh Hoá	Msc. Nghiem Duc Trong	01	HNIP/18257/16
9	TD_09	Le Thanh Binh	05/2015	08/08/2016	<i>Blumea lanceolaria</i> (Roxb.) Druce	Xương sông	Thanh Hoá	Msc. Nghiem Duc Trong	01	HNIP/18258/16
10	TD_10	Le Thanh Binh	05/2015	08/08/2016	<i>Eucalyptus camaldulensis</i> Dehnh.	Bạch đàn trắng	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18259/16
11	TD_11	Le Thanh Binh	2015	08/08/2016	<i>Litsea cubeba</i> (Lour.) Pers.	Màng tang	Quảng Nam	Msc. Nghiem Duc Trong	01	HNIP/18260/16

No.	Sign	Collector & Sender	Date of collection	Date sent	Scientific name	Vietnamese name	Locality	Determined by	Duplicate	Number sign in HNIP
12	TD_12	Le Thanh Binh	06/2015	08/08/2016	<i>Kaempferia galanga</i> L.	Địa liền	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18261/16
13	TD_13	Le Thanh Binh	07/2015	08/08/2016	<i>Zingiber zerumbet</i> (L.) Roscoe ex Sm.	Gừng gió	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18262/16
14	TD_14	Le Thanh Binh	08/2015	08/08/2016	<i>Ageratum conyzoides</i> (L.) L.	Cây cút lợn	Thanh Hoá	Msc. Nghiem Duc Trong	01	HNIP/18263/16
15	TD_18	Le Thanh Binh	07/2015	08/08/2016	<i>Curcuma longa</i> L.	Nghệ	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18264/16
16	TD_19	Le Thanh Binh	07/2015	08/08/2016	<i>Pogostemon cablin</i> (Blanco) Benth.	Hoắc hương	Hung Yên	Msc. Nghiem Duc Trong	01	HNIP/18265/16
17	TD_20	Le Thanh Binh	07/2015	08/08/2016	<i>Hyptis suaveolens</i> (L.) Poit.	É lớn trồng	Vĩnh Phúc	Msc. Nghiem Duc Trong	01	HNIP/18266/16
18	TD_22	Le Thanh Binh	05/2015	08/08/2016	<i>Vitex trifolia</i> L.	Mạn kinh	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18267/16
19	TD_23	Le Thanh Binh	08/2015	08/08/2016	<i>Clausena indica</i> (Dalzell) Oliv.	Mác mật	Thái Nguyên	Msc. Nghiem Duc Trong	01	HNIP/18268/16
20	TD_24	Le Thanh Binh	08/2015	08/08/2016	<i>Elsholtzia blanda</i> (Benth.) Benth.	Kinh giới núi	Lào Cai	Msc. Nghiem Duc Trong	01	HNIP/18269/16
21	TD_25	Le Thanh Binh	08/2015	08/08/2016	<i>Elsholtzia penduliflora</i> W. W. Sm.	Chùa dù	Lào Cai	Msc. Nghiem Duc Trong	01	HNIP/18270/16
22	TD_26	Le Thanh Binh	08/2015	08/08/2016	<i>Litsea cubeba</i> (Lour.) Pers.	Màng tang	Lào Cai	Msc. Nghiem Duc Trong	01	HNIP/18271/16
23	TD_27	Le Thanh Binh	2015	08/08/2016	<i>Illicium verum</i> Hook. f.	Hôi	Lạng Sơn	Msc. Nghiem Duc Trong	01	HNIP/18272/16
24	TD_28	Le Thanh Binh	07/2015	08/08/2016	<i>Zingiber officinale</i> Roscoe	Gừng	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18273/16
25	TD_29	Le Thanh Binh	08/2015	08/08/2016	<i>Hedychium coronarium</i> J.Koenig	Ngải tiên trắng, Bạch yến	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18274/16
26	TD_30	Le Thanh Binh	07/2015	08/08/2016	<i>Coriandrum sativum</i> L.	Mùi	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18275/16
27	TD_32	Le Thanh Binh	07/2015	08/08/2016	<i>Anethum graveolens</i> L.	Thì là	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18276/16

No.	Sign	Collector & Sender	Date of collection	Date sent	Scientific name	Vietnamese name	Locality	Determined by	Duplicate	Number sign in HNIP
28	TD_33	Le Thanh Binh	07/2015	08/08/2016	<i>Plectranthus amboinicus</i> (Lour.) Spreng.	Húng chanh	Phú Thọ	Msc. Nghiem Duc Trong	01	HNIP/18277/16
29	TD_34	Le Thanh Binh	08/2015	08/08/2016	<i>Elsholtzia ciliata</i> (Thunb.) Hyl. (Syn. = <i>Elsholtzia cristata</i> Willd.)	Kinh giới	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18278/16
30	TD_35	Le Thanh Binh	08/2015	08/08/2016	<i>Zingiber montanum</i> (J.Koenig) Link ex A.Dietr.	Gừng núi	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18279/16
31	TD_36	Le Thanh Binh	08/2015	08/08/2016	<i>Amomum schmidtii</i> (K.Schum.) Gagnep.	Sa nhân hồi	Quảng Ninh	Msc. Nghiem Duc Trong	01	HNIP/18280/16
32	TD_38	Le Thanh Binh	11/2014	08/08/2016	<i>Elsholtzia communis</i> (Collett & Hemsl.) Diels	Kinh giới Đồng Văn	Hà Giang	Msc. Nghiem Duc Trong	01	HNIP/18281/16
33	TD_39	Le Thanh Binh	08/2015	08/08/2016	<i>Amomum aromaticum</i> Roxb.	Thảo quả	Lào Cai	Msc. Nghiem Duc Trong	01	HNIP/18282/16
34	TD_40	Le Thanh Binh	08/2015	08/08/2016	<i>Cinnamomum cassia</i> (L.) J.Presl	Quế	Quảng Ninh	Msc. Nghiem Duc Trong	01	HNIP/18283/16
35	TD_42	Le Thanh Binh	08/2015	08/08/2016	<i>Melaleuca cajuputi</i> Powell	Tràm lá dài	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18284/16
36	TD_43	Le Thanh Binh	07/2015	08/08/2016	<i>Alpinia galanga</i> (L.) Willd.	Riềng	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18285/16
37	TD_44	Le Thanh Binh	08/2015	08/08/2016	<i>Curcuma zedoaria</i> (Christm.) Roscoe	Nghệ đen	Hà Nội	Msc. Nghiem Duc Trong	01	HNIP/18286/16

Hanoi, 8th September, 2016

Sender



Le Thanh Binh

Herbarium Keeper



Nghiem Duc Trong



HANOI UNIVERSITY OF PHARMACY, VIETNAM

DEPARTMENT OF BOTANY

IDENTIFICATION OF SCIENTIFIC NAME OF PLANTS

Number: 30/2016

No.	Sign	Collector & Sender	Date of collection	Scientific name	Family	Vietnamese name	Locality
1	TD_01	Le Thanh Binh	05/2015	<i>Melaleuca alternifolia</i> (Maiden & Betch) Cheel	Myrtaceae	Tràm Úc	Hà Nội
2	TD_02	Le Thanh Binh	05/2015	<i>Ocimum gratissimum</i> L.	Lamiaceae	Hương nhu trắng	Hà Nội
3	TD_03	Le Thanh Binh	05/2015	<i>Pluchea indica</i> (L.) Less.	Asteraceae	Cúc tần	Hà Nội
4	TD_04	Le Thanh Binh	05/2015	<i>Platyclusus orientalis</i> (L.) Franco	Cupressaceae	Trắc bách	Hà Nội
5	TD_05	Le Thanh Binh	06/2015	<i>Piper sarmentosum</i> Roxb.	Piperaceae	Lá lốt	Hà Nội
6	TD_06	Le Thanh Binh	08/2015	<i>Artemisia annua</i> L.	Asteraceae	Thanh hao hoa vàng	Hà Nội
7	TD_07	Le Thanh Binh	05/2015	<i>Ocimum tenuiflorum</i> L.	Lamiaceae	Hương nhu tía	Hà Nội
8	TD_08	Le Thanh Binh	08/2015	<i>Dysphania ambrosioides</i> (L.) Mosyakin & Clemant	Chenopodiaceae	Dầu giun	Thanh Hoá
9	TD_09	Le Thanh Binh	05/2015	<i>Blumea lanceolaria</i> (Roxb.) Druce	Asteraceae	Xương sông	Thanh Hoá
10	TD_10	Le Thanh Binh	05/2015	<i>Eucalyptus camaldulensis</i> Dehnh.	Myrtaceae	Bạch đàn trắng	Hà Nội
11	TD_11	Le Thanh Binh	2015	<i>Litsea cubeba</i> (Lour.) Pers.	Lauraceae	Màng tang	Quảng Nam
12	TD_12	Le Thanh Binh	06/2015	<i>Kaempferia galanga</i> L.	Zingiberaceae	Địa liên	Hà Nội
13	TD_13	Le Thanh Binh	07/2015	<i>Zingiber zerumbet</i> (L.) Roscoe ex Sm.	Zingiberaceae	Gừng gió	Hà Nội
14	TD_14	Le Thanh Binh	08/2015	<i>Ageratum conyzoides</i> (L.) L.	Asteraceae	Cây cứt lợn	Thanh Hoá
15	TD_18	Le Thanh Binh	07/2015	<i>Curcuma longa</i> L.	Zingiberaceae	Nghệ	Hà Nội

No.	Sign	Collector & Sender	Date of collection	Scientific name	Family	Vietnamese name	Locality
16	TD_19	Le Thanh Binh	07/2015	<i>Pogostemon cablin</i> (Blanco) Benth.	Lamiaceae	Hoặc hương	Hung Yên
17	TD_20	Le Thanh Binh	07/2015	<i>Hyptis suaveolens</i> (L.) Poit.	Lamiaceae	É lớn trồng	Vĩnh Phúc
18	TD_22	Le Thanh Binh	05/2015	<i>Vitex trifolia</i> L.	Verbenaceae	Mạn kinh	Hà Nội
19	TD_23	Le Thanh Binh	08/2015	<i>Clausena indica</i> (Dalzell) Oliv.	Rutaceae	Mác mật	Thái Nguyên
20	TD_24	Le Thanh Binh	08/2015	<i>Elsholtzia blanda</i> (Benth.) Benth.	Lamiaceae	Kinh giới núi	Lào Cai
21	TD_25	Le Thanh Binh	08/2015	<i>Elsholtzia penduliflora</i> W. W. Sm.	Lamiaceae	Chùa dù	Lào Cai
22	TD_26	Le Thanh Binh	08/2015	<i>Litsea cubeba</i> (Lour.) Pers.	Lauraceae	Màng tang	Lào Cai
23	TD_27	Le Thanh Binh	2015	<i>Illicium verum</i> Hook. f.	Illiciaceae	Hôi	Lạng Sơn
24	TD_28	Le Thanh Binh	07/2015	<i>Zingiber officinale</i> Roscoe	Zingiberaceae	Gừng	Hà Nội
25	TD_29	Le Thanh Binh	08/2015	<i>Hedychium coronarium</i> J.Koenig	Zingiberaceae	Ngải tiên trắng, Bạch yến	Hà Nội
26	TD_30	Le Thanh Binh	07/2015	<i>Coriandrum sativum</i> L.	Apiaceae	Mùi	Hà Nội
27	TD_32	Le Thanh Binh	07/2015	<i>Anethum graveolens</i> L.	Apiaceae	Thì là	Hà Nội
28	TD_33	Le Thanh Binh	07/2015	<i>Plectranthus amboinicus</i> (Lour.) Spreng.	Lamiaceae	Húng chanh	Phú Thọ
29	TD_34	Le Thanh Binh	08/2015	<i>Elsholtzia ciliata</i> (Thunb.) Hyl. (Syn. = <i>Elsholtzia cristata</i> Willd.)	Lamiaceae	Kinh giới	Hà Nội
30	TD_35	Le Thanh Binh	08/2015	<i>Zingiber montanum</i> (J.Koenig) Link ex A.Dietr.	Zingiberaceae	Gừng núi	Hà Nội
31	TD_36	Le Thanh Binh	08/2015	<i>Amomum schmidtii</i> (K.Schum.) Gagnep.	Zingiberaceae	Sa nhân hôi	Quảng Ninh
32	TD_38	Le Thanh Binh	11/2014	<i>Elsholtzia communis</i> (Collett & Hemsl.) Diels	Lamiaceae	Kinh giới Đông Văn	Hà Giang
33	TD_39	Le Thanh Binh	08/2015	<i>Amomum aromaticum</i> Roxb.	Zingiberaceae	Thảo quả	Lào Cai

No.	Sign	Collector & Sender	Date of collection	Scientific name	Family	Vietnamese name	Locality
34	TD_40	Le Thanh Binh	08/2015	<i>Cinnamomum cassia</i> (L.) J.Presl	Lauraceae	Quế	Quảng Ninh
35	TD_42	Le Thanh Binh	08/2015	<i>Melaleuca cajuputi</i> Powell	Myrtaceae	Tràm lá dài	Hà Nội
36	TD_43	Le Thanh Binh	07/2015	<i>Alpinia galanga</i> (L.) Willd.	Zingiberaceae	Riềng	Hà Nội
37	TD_44	Le Thanh Binh	08/2015	<i>Curcuma zedoaria</i> (Christm.) Roscoe	Zingiberaceae	Nghệ đen	Hà Nội

Hanoi, 8th September, 2016

Head of Department

Professor Tran Van On

Determiner

Msc. Nghiem Duc Trong

References

1. Delin Wu, Kai Larsen (2000), Zingiberaceae, in: Wu, Z. Y. & P. H. Raven (eds.), *Flora of China, Vol. 24 (Flagellariaceae through Marantaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
2. Dianxiang Zhang, Thomas G. Hartley, David J. Mabberley (2008), Rutaceae, in: Wu, Z. Y., P. H. Raven & D. Y. Hong (eds.), *Flora of China. Vol. 11 (Oxalidaceae through Aceraceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
3. Gelin Zhu et al. (2003), Chenopodiaceae, in: Wu, Z. Y., P. H. Raven & D. Y. Hong (eds.), *Flora of China, Vol. 5 (Ulmaceae through Basellaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
4. Jie Chen, Lyn A. Craven (2007), Myrtaceae, in: Wu, Z. Y., P. H. Raven & D. Y. Hong (eds.), *Flora of China, Vol. 13 (Chusiaceae through Araliaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
5. Le Kim Bien (2007) Asteraceae in: Nguyen Tien Ban et al (eds.), *Flora of Vietnam, Volume 7*, Science and Technics Publishing House, Hanoi
6. Ligu Fu et al. (1999), Cupressaceae, in: Wu, Z. Y. & P. H. Raven (eds.), *Flora of China, Vol. 4 (Cycadaceae through*

Fagaceae), Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.

7. Menglan She et al. (2005), Apiaceae, in: Wu, Z. Y., P. H. Raven & D. Y. Hong (eds.), *Flora of China, Vol. 14 (Apiaceae through Ericaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
8. Nianhe Xia, Richard M. K. Saunders (2008), Illiciaceae, in: Wu, Z. Y., P. H. Raven & D. Y. Hong (eds.), *Flora of China, Vol. 7 (Menispermaceae through Capparaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
9. Pham Hoang Ho (1999-2000), *Vietnam Plants (An Illustrated Flora of Vietnam), Volume 1-3*, Youth Publishing House, Ho Chi Minh City
10. Shou-liang Chen, Michael G. Gilbert (1994), Verbenaceae, in: Wu, Z. Y. & P. H. Raven (eds.), *Flora of China, Vol. 17 (Verbenaceae through Solanaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
11. Shugang Li et al. (2008), Lauraceae, in: Wu, Z. Y., P. H. Raven & D. Y. Hong, (eds.), *Flora of China, Vol. 7 (Menispermaceae through Capparaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
12. Vo Van Chi (2004), *Dictionary of common plants, Volume 1-2*, Science and Technics Publishing House, Hanoi
13. Vu Xuan Phuong (2000), Lamiaceae, in: Nguyen Tien Ban et al (eds.), *Flora of Vietnam, Volume 2*, Science and Technics Publishing House, Hanoi
14. Vu Xuan Phuong (2007), Verbenaceae, in: Nguyen Tien Ban et al (eds.), *Flora of Vietnam, Volume 6*, Science and Technics Publishing House, Hanoi
15. Xi-wen Li, Ian C. Hedge (1994), Lamiaceae, in: Wu, Z. Y. & P. H. Raven (eds.), *Flora of China, Vol. 17 (Verbenaceae through Solanaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
16. Yung-chien Tseng, Nianhe Xia, Michael G. Gilbert (1999), Piperaceae, in: Wu, Z. Y. & P. H. Raven (eds.), *Flora of China, Vol. 4 (Cycadaceae through Fagaceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
17. Zhu Shi et al. (2011), Asteraceae (Compositae), in: Wu, Z. Y., Raven, P. H. & Hong, D. Y. (eds.), *Flora of China, Volume 20-21 (Asteraceae)*, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.