



The Persister Character of Clinical Isolates of *Staphylococcus aureus* Contributes to Faster Evolution to Resistance and Higher Survival in THP-1 Monocytes: A Study With Moxifloxacin

Tiep K. Nguyen^{1,2}, Frédéric Peyrusson¹, Magali Dodémont³, Nhung H. Pham^{4,5}, Hoang A. Nguyen⁶, Paul M. Tulkens¹ and Françoise Van Bambeke^{1*}

¹ Pharmacologie Cellulaire et Moléculaire, Louvain Drug Research Institute, Université catholique de Louvain (UCLouvain), Brussels, Belgium, ² Department of Pharmaceutical Industry, Hanoi University of Pharmacy, Hanoi, Vietnam, ³ Centre National de Référence des Staphylocoques, Laboratoire Hospitalier Universitaire de Bruxelles (LHUB-ULB) Site Anderlecht, Hôpital Erasme – Cliniques Universitaires de Bruxelles, Brussels, Belgium, ⁴ Department of Microbiology, Bach Mai Hospital, Hanoi, Vietnam, ⁵ Microbiology Department, Hanoi Medical University, Hanoi, Vietnam, ⁶ The National Center for Drug Information and Adverse Drug Reactions Monitoring, Hanoi University of Pharmacy, Hanoi, Vietnam

OPEN ACCESS

Edited by:

Miklos Fuzi,
Semmelweis University, Hungary

Reviewed by:

Niels Frimodt-Møller,
Rigshospitalet, Denmark
Yu Zhijian,
Shenzhen University, China

*Correspondence:

Françoise Van Bambeke
francoise.vanbambeke@uclouvain.be

Specialty section:

This article was submitted to
Antimicrobials, Resistance
and Chemotherapy,
a section of the journal
Frontiers in Microbiology

Received: 25 July 2020

Accepted: 16 October 2020

Published: 23 November 2020

Citation:

Nguyen TK, Peyrusson F, Dodémont M, Pham NH, Nguyen HA, Tulkens PM and Van Bambeke F (2020) The Persister Character of Clinical Isolates of *Staphylococcus aureus* Contributes to Faster Evolution to Resistance and Higher Survival in THP-1 Monocytes: A Study With Moxifloxacin. *Front. Microbiol.* 11:587364. doi: 10.3389/fmicb.2020.587364

Staphylococcus aureus may cause relapsing infections. We previously showed that *S. aureus* SH1000 surviving intracellularly to bactericidal antibiotics are persisters. Here, we used 54 non-duplicate clinical isolates to assess links between persistence, resistance evolution, and intracellular survival, using moxifloxacin throughout as test bactericidal antibiotic. The relative persister fraction (RPF: percentage of inoculum surviving to 100× MIC moxifloxacin in stationary phase culture for each isolate relative to ATCC 25923) was determined to categorize isolates with low (≤ 10) or high (> 10) RPF. Evolution to resistance (moxifloxacin MIC ≥ 0.5 mg/L) was triggered by serial passages at 0.5× MIC (with daily concentration readjustments). Intracellular moxifloxacin maximal efficacy (E_{\max}) was determined by 24 h concentration-response experiments [pharmacodynamic model (Hill-Langmuir)] with infected THP-1 monocytes exposed to moxifloxacin (0.01 to 100× MIC) after phagocytosis. Division of intracellular survivors was followed by green fluorescence protein dilution (FACS). Most (30/36) moxifloxacin-susceptible isolates showed low RPF but all moxifloxacin-resistant ($n = 18$) isolates harbored high RPF. Evolution to resistance of susceptible isolates was faster for those with high vs. low RPF (with SOS response and topoisomerase-encoding genes overexpression). Intracellularly, moxifloxacin E_{\max} was decreased (less negative) for isolates with high vs. low RPF, independently from resistance. Moxifloxacin intracellular survivors were non-dividing. The data demonstrate and quantitate persisters in clinical isolates of *S. aureus*, and show that this phenotype accelerates resistance evolution and is associated with intracellular survival in spite of high antibiotic concentrations. Isolates with high RPF may represent a possible cause of treatment failure not directly related to resistance in patients receiving active antibiotics.

Keywords: *Staphylococcus aureus*, persister, intracellular infection, resistance, intracellular survival, moxifloxacin

INTRODUCTION

Persisters, defined as subpopulations of bacteria that do not multiply but still survive in the presence of a bactericidal antibiotic, are suspected of being a potential cause of therapeutic failure in spite of the use of active antibiotics (Van den Bergh et al., 2017; Jung et al., 2019). This phenotype is not genetically inherited and is fully reversible upon antibiotic removal (Cohen et al., 2013), thereby escaping detection through conventional antibiotic susceptibility testing. Persisters grow normally in conventional media, and need, therefore, to be distinguished from small colony variants (SCVs), which are phenotypic subpopulations harboring specific auxotrophisms leading to slow growth and small colonies formation; SCVs also easily revert to the original wild-type phenotype depending on environmental conditions and are sometimes persisters (Tuchscherr et al., 2020). Antibiotic persistence is also distinct from resistance, characterized by an increase in the minimal inhibitory concentration (MIC) but maintenance of the bacterial capacity to multiply as long as the antibiotic concentration is kept below its current MIC. Moreover, resistance phenotypes are often genetically inherited and poorly reversible upon antibiotic withdrawal. Yet, it has been proposed that persisters may constitute an evolutionary reservoir from which resistance can emerge (Cohen et al., 2013; Windels et al., 2019). Conversely, tolerance, characterized by a reduced killing rate of bacteria (as also observed in persisters), can interact synergistically with resistance to help bacteria escaping antibiotic killing by combining these strategies (Ortiz-Padilla et al., 2020).

Persisters are also associated with specific modes of life in which bacteria show apparent unresponsiveness to antibiotics as long as they are maintained in the corresponding specific environment. Focusing on intracellular infection, the presence of persisters within the host cells may be a key determinant in the intracellular survival of small but significant inocula for many bacteria (Fisher et al., 2017). We and others have further documented that antibiotics are unable to eradicate intracellular *Staphylococcus aureus* even upon exposure of the host cells to high concentrations (typically 100-fold their MIC) of highly bactericidal agents (Krut et al., 2004; Barcia-Macay et al., 2006; Lemaire et al., 2011; Peyrusson et al., 2019), leaving apparently untouched about 0.1–1% of the original inoculum (Peyrusson et al., 2019). This contrasts with the observed extracellular activity of these drugs, which are capable of bringing CFU counts below the limit of detection in a very short time and often at concentrations that are only a few multiples of their MIC. Using the *S. aureus* laboratory strain SH1000, we recently showed that intracellular survivors collected from macrophages or monocytes exposed to high antibiotic concentrations are actually persisters (Peyrusson et al., 2020). This conclusion was supported by the demonstration of a non-dividing state for these survivors as long as they remained intracellular and under antibiotic pressure, but a rapid reversion to a growth state as soon as the antibiotic pressure was relieved; it was accompanied by an in-depth transcriptomic analysis revealing an activation of the stringent response, the cell wall stress stimulon, and the SOS and heat shock responses,

together with profound changes in metabolic fluxes. However, the occurrence and quantification of persisters in clinical isolates, as well as the impact of their relative abundance on (i) evolution to resistance of the whole population and (ii) lack of eradication from phagocytes exposed to large concentrations of an active antibiotic have so far not been established.

In an attempt to address these unanswered questions, we assembled 36 non-duplicated isolates from patients hospitalized in Hanoi, Vietnam, and presenting infections that did not resolve or reactivated after 5 days of treatment with an active antibiotic (Nguyen et al., 2020), complemented with 18 randomly selected clinical isolates from the Belgian reference center for Staphylococci to give a more generalizable character to this collection. We first looked for the relative abundance of persisters in stationary cultures of these isolates, using the fluoroquinolone moxifloxacin as selecting agent based on its highly bactericidal activity against susceptible *S. aureus* in broth, and as being also one of the most active antibiotics against intracellular susceptible *S. aureus* invading and thriving in human permissive monocytes (Barcia-Macay et al., 2006; Lemaire et al., 2011; Peyrusson et al., 2020). We then compared isolates with low vs. high persisters abundance (referred henceforth as isolates with a high- and low-persister character, respectively) (i) for evolution to resistance to moxifloxacin (MIC increase) upon serial exposure to subinhibitory concentrations of this antibiotic in broth (a model previously developed and applied to the study of evolution of pneumococcal resistance to fluoroquinolones (Avrain et al., 2007), and (ii) for determination of the eradication of phagocytized bacteria from permissive monocytes exposed to moxifloxacin [using a suitable pharmacodynamic approach (Buyck et al., 2016)], together with an analysis of the capacity of survivors to divide intracellularly. All moxifloxacin-resistant isolates in the collection had a high-persister character while most but not all susceptible isolates showed a low-persister character. Most interestingly, susceptible isolates with a high-persister character evolved more rapidly to resistance in broth than those with a low-persister character. They were also lesser efficiently eradicated from phagocytes exposed to moxifloxacin (independently of their resistance pattern), with all intracellular survivors showing evidence of minimal division. Collectively, the data suggest that a high-persister character in a clinical isolate of *S. aureus* may be a contributing factor to therapeutic failures (Garzoni and Kelley, 2009; Conlon, 2014; Dubourg et al., 2017; Boudjemaa et al., 2018).

MATERIALS AND METHODS

Main Products

Moxifloxacin HCl (potency: 90.9%) was obtained from Bayer HealthCare (Leverkusen, Germany); gentamicin sulfate (potency: 60.7%) and cation-adjusted Muller Hinton broth (CA-MHB), from Sigma-Aldrich (St. Louis, MO, United States); human serum, from Biowest SAS (Nuaillé, France); cell culture media, from Gibco/Life Technologies Corporation (Paisley, United Kingdom); Mueller Hinton broth (MHB) and tryptic soy

agar (TSA), from VWR (Radnor, PA, United States); and primers, from Eurogentec (Liège, Belgium).

Bacterial Strains and Antibiotic Susceptibility Testing

Staphylococcus aureus ATCC 25923 and RN 4220 are reference strains from the American Tissue Culture Collection (ATCC), Manassas, VA, United States. Thirty-six clinical isolates were collected at the Bach Mai Hospital (Hanoi, Vietnam) from patients with infections that did not resolve after 5 days of treatment by antibiotics to which the initial isolates were reported as susceptible or that reactivated after treatment discontinuation (Nguyen et al., 2020). Eighteen isolates from the collection of the Belgian National Reference Centre for Staphylococci were randomly selected and included to give a more general character to our observations. **Supplementary Table 1** shows, for all these strains, their MSSA/MRSA phenotype, the MIC of moxifloxacin [determined by broth microdilution (according to the Clinical and Laboratory Standards Institute recommendations)], the moxifloxacin relative persister fraction (RPF; see next paragraph), and the corresponding assigned phenotype.

Determination and Quantification of Persistence *in vitro*

Bacteria were grown to stationary phase after 2 sub-cultures in CA-MHB and then exposed to moxifloxacin at 100× MIC or its vehicle (sterile water; control) in CA-MHB for 5 h at 37°C under gentle rotary shaking (130 rpm). CFUs were counted after plating and overnight incubation of serially diluted aliquots. The percentage of persisters was calculated as the ratio of the CFU number in each moxifloxacin-treated sample to that in the corresponding control, and the RPF, as the ratio of the percentage of persisters for each clinical isolate to that observed for ATCC 25923 (De Groote et al., 2009).

In vitro Selection of Resistance

Bacteria from an overnight broth culture were inoculated in 12-well plates (0.5×10^6 CFU/mL) in CA-MHB and exposed to moxifloxacin at 0.5× MIC, with daily measurement of the MIC and readjustment of the moxifloxacin concentration to half of this value until the MIC reached a value of 2 mg/L (Avrain et al., 2007). This experimental set-up referred to as “serial passage” is widely used to follow evolution toward resistance (Martinez et al., 2007) especially when it is related to mutations in target genes or gene regulators. It does not mimic what could occur in clinical practice when patients are exposed to optimal dosing regimens, but rather in case of antibiotic underdosing known as a main driver of evolution toward resistance (Heffernan et al., 2018).

Determination of Gene Expression Levels by RT-PCR

Stationary-phase cultures were incubated with moxifloxacin or its vehicle as for the persister assay, and harvested by centrifugation either immediately or after incubation with moxifloxacin at 100× MIC. Bacteria were lysed with lysozyme and lysostaphin, after which RNA was isolated (InviTrap® Spin Universal RNA Mini

Kit; Stratec, Berlin, Germany) and purified (TURBO DNA-free™ Kit; Invitrogen, Carlsbad, CA, United States). The RNA quality and quantity were determined by spectrophotometry (NanoDrop™, Thermofisher, Waltham, MA, United States). Purified RNA was converted to cDNA (Transcription first strand cDNA synthesis kit; Roche Applied Science, Penzberg, Germany). Quantitative RT-PCR was performed with a iCycler™ iQ instrument (Bio-Rad, Hercules, CA, United States) using 5 µL of cDNA, 12.5 µL of SYBR Green Master Mix (Bio-Rad), 2 µL of 5 mM of each primer and 3.5 µL of sterile RNase-free water (Ambion, St. Austin, TX, United States). PCR program steps were (i) 95°C 3 min (denaturation) (ii) 40 cycles (95°C 15 s–61°C 60 s) (amplification). Primers are described in **Supplementary Table 2**. A melting curve was run to check for the presence of a unique PCR reaction product. Relative expression levels were calculated by the $\Delta\Delta C_t$ method, with *gmk* as housekeeping gene and *S. aureus* ATCC 25923 as a reference.

Screening for Mutations in the Genes Encoding Fluoroquinolone Targets

Template DNA was purified with GeneJET Genomic DNA Purification Kit (Thermofisher). PCR amplification of the QRDRs of *gyrA*, *gyrB*, *parC*, and *parE* was performed using 1 µL of DNAG, 5 µL of 5× Phusion buffer, 0.8 µL of 10 mM dNTPs, 1 µL of 10 mM of each primer (see **Supplementary Table 2**), 0.25 µL of Phusion polymerase (Thermofisher) and 15.95 µL sterile nuclease-free water (Ambion). The PCR program steps were (i) 98°C 3 min (denaturation), (ii) 30 cycles [98°C 10 s–60°C 30 s (56°C 30s with *parE*), 72°C 60 s] (amplification), and (iii) 72°C 10 min (full amplicons extension). Sanger sequencing was performed in forward and reverse directions at Genewiz (Leipzig, Germany) using the same primers. QRDR DNA sequences were compared with those of *S. aureus* RN4220.

Intracellular Activities of Antibiotics

THP-1 human monocytes (clone ATCC TIB-202; American Type Culture Collection) were infected with *S. aureus* as previously described, and non-phagocytized bacteria eliminated by short exposure (45 min) to gentamicin at 50 to 100× its MIC (thus limiting the study to isolates for which the MIC of gentamicin was ≤ 8 mg/L (i.e., a maximum of 4-fold the current EUCAST cut-off of 2 mg/L observed for wild type strains) (Barcia-Macay et al., 2006). Results are expressed as the change in the number of CFU per mg of cell protein after 24 h incubation of infected cells with moxifloxacin. We checked in preliminary experiments that moxifloxacin, including at high concentrations, did not affect the protein content of the samples neither the number of cells after 24 h of incubation. We also checked that proteins from dead cells do not interfere in the protein assay (signal barely distinguishable from that of the blank for cells killing by incubation with DMSO or Triton X-100). The activity of moxifloxacin against intracellular bacteria was measured using full concentration-response experiments (change in intracellular CFU from the post-phagocytosis inoculum plotted against the extracellular antibiotic concentration [both in log₁₀ units] typically ranging from 10^{−2} to 10⁺² fold the MIC). A Hill-Langmuir equation

[with slope factor set to 1 as discussed previously (Lemaire et al., 2009; Peyrusson et al., 2019)] was fitted to the data to calculate two key pharmacodynamic parameters [apparent static concentration (C_s) and maximal relative efficacy (E_{max}); see Buyck et al. (2016) and in the caption of **Figure 4**].

Transformation of *Staphylococcus aureus*

gfp, encoding the Green fluorescent protein (GFP), cloned with the *xyl/tetO* inducible promoter carried on a shuttle pALC 2084 *Escherichia coli* – *S. aureus* plasmid was used as a reporter (Bateman et al., 2001) transformed into *S. aureus* by electroporation following exactly a previously described protocol (Rhoads Kraemer and Iandolo, 1990). Colonies were checked for GFP expression (fluorescence microscopy) after incubation with 0.125 mg/L tetracycline (inducer), and for unimpaired susceptibility to moxifloxacin, gentamicin, and chloramphenicol.

Flow Cytometry

Transformed bacteria were cultured overnight in CA-MHB supplemented with 0.125 mg/L tetracycline (GFP inducer) and 10 mg/L chloramphenicol. They were used for intracellular infection following our general protocol, except that the phagocytosis time was reduced to 30 min (to limit bacterial division during this step), and the ratio of bacteria to THP-1 cells doubled (to ease the fluorometric quantitation of the intracellular inoculum). Tetracycline was removed from the medium at the end of the phagocytosis step. CFUs counting and FACS analysis were performed in parallel on the same samples as previously described (Peyrusson et al., 2020). In brief, bacteria isolated from THP-1 cells were resuspended in filtered PBS, stained with 10 mg/L propidium iodide, and analyzed using a FACSVerse cytometer (BD Biosciences, Franklin Lakes, NJ, United States) for GFP signal intensities (FITC channel, medium flow rate). Forward-scatter width (FCS-W) vs. forward-scatter area (FSC-A), and side-scatter width (SSC-W) vs. side-scatter area (SSC-A) were used to gate out damaged (propidium iodide-positive) or multiplet bacteria (Peyrusson et al., 2020). Data were analyzed with FlowJo 10.5.2 software (TreeStar Inc., Ashland, OR, United States).

RESULTS

Determination of the Moxifloxacin-Persister Character of Clinical Isolates

Figure 1A shows the killing of ATCC 25923 and 2 selected Vietnamese clinical isolates in stationary phase cultures over time of exposure to moxifloxacin at 100× MIC. A plateau of killing [typical of persisters (Balaban et al., 2019)] was reached after about 5 h. **Figure 1B** shows the RPF for all Vietnamese and Belgian isolates, categorized according to their susceptibility to moxifloxacin (individual values in **Supplementary Table 1**). Among moxifloxacin-susceptible isolates, we could categorize the isolates as having a low (≤ 10 ; $n = 30/36$) or a high (> 10 ; $n = 6/36$)

RPF. All moxifloxacin-resistant isolates ($n = 18$) showed a high RPF. We therefore considered three phenotypic subclasses for this work, henceforth referred to as susceptible isolates with low relative persister fractions (S-LP), or with high relative persister fractions (S-HP), and resistant isolates with high relative persister fractions (R-HP), respectively¹.


Expression of Genes Involved in SOS Response and Encoding Fluoroquinolone Targets

Exposure of bacteria to fluoroquinolones triggers a SOS response, which, in turn, increases mutation frequency (Qin et al., 2015) and development of resistance, but also induces persistence (Dorr et al., 2009). Using representative isolates of the three phenotypic groups defined above and exposed to moxifloxacin as for the persister assay, we measured the expression of the SOS response regulon repressor *lexA*, the derepressor *recA*, and two genes of this regulon [*uvrB*, encoding the excision nuclease subunit B recognizing DNA lesions and involved in repair processes initiated by fluoroquinolone-induced damage; *umuC*, encoding an error-prone polymerase introducing mutations in DNA during replication (Qin et al., 2015; Maslowska et al., 2019)]. The expression of these genes was induced in all isolates, except for *recA* (overexpressed in S-LP, to a lesser extent in S-HP, but not in R-HP; **Figure 2A**). Because genes encoding fluoroquinolone targets are potentially regulated by the SOS response (Cirz et al., 2007), we measured their expression in the same conditions (**Figure 2B**). We observed an overexpression in S-HP and R-HP but not in S-LP.

Selection of Resistance to Moxifloxacin in Susceptible Isolates With Low- and High-Relative Persister Fractions

Since all moxifloxacin-resistant isolates showed high relative persister fractions, we tested whether S-HP isolates would be more prone to develop resistance than their S-LP counterparts. We used a previously developed model (Avrain et al., 2007) in which bacteria are exposed to subinhibitory concentrations of a fluoroquinolone in a closed system (excluding therefore the acquisition of foreign genetic material), to specifically assess the intrinsic ability of isolates for spontaneous evolution to develop resistance while avoiding selection of pre-existing subpopulations with higher MIC. Thus, 3 S-LP and 3 S-HP isolates were exposed to 0.5× MIC of moxifloxacin and the change in MIC measured (and moxifloxacin concentration

¹To avoid any ambiguity, S or R in the abbreviation of susceptible or resistant isolates refer to the MIC measured in broth and consider therefore a property attributed to (or assumed to be shared by) each individual organism in the entire population (thus, disregarding potential differences in MIC among individual organisms that could be revealed, for instance, by diffusion or population analysis techniques). Conversely, the “HP” or “LP” abbreviations used to denote isolates with a high- or a low-persister character refer to a low or high percentage of bacteria isolated from the initial population relative to the value observed with the laboratory strain ATCC 25923. It, therefore, does not mean that the persister character concerns the whole population, since actually, most of the individual bacteria are not persisters. It rather refers to a small subpopulation and describes its relative abundance in the whole population (selected during the assay) in the conditions of the assay.




readjusted) each day. **Figure 3A** (with data limited to the first 15 days of the experiment) shows that resistance was acquired faster for S-HP than for S-LP [mean time for onefold MIC increase, using data from day 2 through 15: 2.44 ± 0.37 vs. 4.34 ± 0.77 days ($n = 2 \times 14$; $p < 0.0001$; two tailed unpaired t -test with Welch's correction for unequal SD's)]. At the end of the selection process, mutants (with an MIC of 2 mg/L) had a RPF of approx. 100 [significantly increased for S-LP only, to levels similar to those observed in R-HP (**Figure 3B**)]. The RPF and moxifloxacin MIC increased in parallel in S-LP during this process (**Figure 3C**), with all strains with a MIC ≥ 0.5 mg/L showing a RPF > 10 (**Figure 3D**). We also evaluated in these mutants the expression of genes involved in SOS response and encoding fluoroquinolone targets. They were all overexpressed to levels similar or even higher than in their parental isolates (**Supplementary Figure 1**).

Mechanisms of Resistance to Moxifloxacin

Supplementary Table 3 shows the mutations observed in the QRDR of *gyrA*, *gyrB*, *parC*, and *parE* of selected isolates

from the three phenotypic groups and of resistant mutants selected *in vitro* from S-LP and S-HP. In most susceptible isolates, we noticed a Glu422Asp substitution in ParE, known as not causing resistance (Schmitz et al., 1998). Isolate 69687, which showed a more elevated moxifloxacin MIC, harbored a His103Tyr mutation in ParC, previously described as frequently associated with other mutations in resistant strains (Morrow et al., 2011). All moxifloxacin-resistant clinical isolates presented a Ser84Leu mutation in GyrA and a Ser80Phe mutation in ParC, which together are known as conferring resistance (Roychoudhury et al., 2001; Kwak et al., 2013) and are described in isolates for which moxifloxacin MICs are in the 1–4 mg/L range (Schmitz et al., 1998; Nakaminami et al., 2014). For moxifloxacin-resistant mutants obtained from S-LP isolates, mutations in GyrA (Ser84Leu), GyrB (Leu433Ile or Ile454Val), or ParC (Ser80Phe or Ser80Ala) were detected. To our knowledge, these mutations in GyrB or the substitution of Ser80Ala in ParC have not been previously described. Thus, each of the final mutants showed two mutations in fluoroquinolone targets, which is what is generally required for moxifloxacin to reach MICs ≥ 1 mg/L (Noguchi et al., 2005; Bhagwat et al., 2006). For moxifloxacin-resistant mutants obtained from S-HP, only one mutation in GyrA (Ser80Leu) was




detected, which was previously found in rare resistant isolates (Kwak et al., 2013).

Genes encoding efflux transporters *norA*, *norB*, and *norC* were slightly overexpressed in H-RP, and *norB*, in *in vitro* selected mutants, after incubation with moxifloxacin at 100× MIC, when compared to ATCC 25923 (Supplementary Figure 2).

Quantitative Measurement of Intracellular Activity of Moxifloxacin

In a next step, we evaluated the intracellular activity of moxifloxacin against isolates representative of the S-LP, S-HP, or R-HP phenotypes [selected based on a gentamicin MIC ≤ 8 mg/L; see methods and Barcia-Macay et al. (2006)]. The post-phagocytosis inoculum was similar among the three phenotypic groups [6.4 ± 0.1 , 6.5 ± 0.2 , and 6.7 ± 0.4 log₁₀ CFU/mg protein for isolates with S-LP, S-HP, and R-HP character, respectively (not statistically different by one-way ANOVA)]. Figures 4A–C shows the moxifloxacin patterns of activity in

concentration-response experiments after 24 h of incubation with equipotent antibiotic concentrations (same multiples of the MIC). A Hill-Langmuir function could be fitted to all data, with a single equation for all isolates within each phenotypic group. The static concentrations (C_s) were close to the corresponding MICs, denoting similar relative potencies. In contrast, the moxifloxacin maximal relative activity [E_{max} , which the model fitted to the data determines by extrapolation of the sigmoid function for an infinitely large antibiotic concentration (describing, therefore, the maximal amplitude of the bacterial response to the antibiotic)], was significantly lower (less negative) for S-HP or R-HP than for S-LP [-1.45 ± 0.31 and -1.33 ± 0.30 vs. -2.10 ± 0.34 log₁₀ CFU (Figure 4D); these values, which correspond to a bottom plateau in the function, are actually close to those measured at the highest extracellular moxifloxacin concentration tested]. An inverse correlation was also observed between the E_{max} value and the RPF of each isolate (Figure 4E). Thus, these data show that cell infection by isolates with high relative persister fractions






FIGURE 4 | Intracellular activity of moxifloxacin. Top (A–C) Concentration-response curves of the activity of moxifloxacin against intraphagocytic *S. aureus* with different resistance/persistence phenotypes [(A): isolates 30337, 69474, 69687, 34467, 2017S228, 2016S354, 2014S118, and 1078 with susceptible low-persister character (S-LP); (B): isolates 69505, 69783, 69867, 36606, and 1214 with susceptible high-persister character (S-HP); (C): isolates 20975, 13890, 2015S418, and 2017S017 with resistant high-persister (R-HP) character]. Infected THP-1 monocytes were incubated for 24 h with the antibiotic at concentrations (expressed in \times MIC of each individual isolate) covering a large span to allow for a full concentration-response analysis (see Buyck et al., 2016). The horizontal dotted line shows the initial post-phagocytosis inoculum. Data are expressed as changes from this value after 24 h of incubation. The dots on each graph correspond to experimental values for individual isolates (with their respective SD) sharing the same resistance-persistence phenotype and the curve is a single Hill-Langmuir function (with slope factor set to 1) fitted to the data of all isolates in each group (with the dotted lines corresponding to its 95% confidence interval; see **Supplementary Table 4** (Continued)

FIGURE 4 | Continued

(Supplementary Material) for individual Hill-Langmuir function parameters of interest [E_{max} (maximal relative efficacy) and IC_{50}]. The vertical line points to the MIC. Middle (D) Maximal relative efficacy (E_{max}) of moxifloxacin (reduction in intracellular CFUs as extrapolated for an infinitely large antibiotic concentration based on the Hill equation parameters determined for each individual strain). The dots are the values of each individual isolate; the horizontal lines, are the mean and SD of each phenotypic group. Statistical analysis: one-way ANOVA ($p = 0.0025$) with Tukey *post hoc* test comparing the three phenotypic subgroups: $**p < 0.01$; there was no significant difference between the S-HP and R-HP groups. (E) Correlation between the intracellular E_{max} (determined as shown here) and the relative persister fraction [as determined in stationary phase culture (see Supplementary Table 1), using log scale for both. The vertical dotted shows the relative persister fraction of the reference ATCC 25923. r and p : Pearson coefficient and p value of the correlation. Bottom (F) Concentration-response curves of the activity of moxifloxacin against intraphagocytic RN4220 (S-LP) and 1214 (S-HP) transformed by the pALC2084 plasmid to express GFP under induction by tetracycline (maintained till the end of the phagocytosis period), as measured after 24 h of incubation of the infected cells in the presence of moxifloxacin over a wide range of concentrations (expressed in \times MIC of each individual isolate) but in the absence of tetracycline. The Hill-Langmuir function (with slope factor set to 1) fitted to the data of each isolate is shown, together with the E_{max} of each curve with its 95% confidence interval at an extrapolated infinitely large concentration. The horizontal dotted line shows the initial post-phagocytosis inoculum; data are expressed as changes from this value after 24 h of incubation. The vertical line points to the MIC. The table above the graph shows the MIC and relative persister fraction of these strains. Yellow rectangles highlight samples that were examined in FACS, with the corresponding concentrations shown by black arrows. (G) Flow cytometric profiles of bacteria recovered from monocytes at the end of the phagocytosis period (0 h) or after 24 h of incubation with moxifloxacin at the indicated concentrations. The graphs show flow cytometric profiles of the frequency of events as a function of GFP fluorescence intensity.

results in larger intracellular residual inocula that remained unaffected by moxifloxacin compared to what is observed for an isolate with a low relative persister fraction. In addition, we found that the growth rate of bacteria collected from intracellular experiments and reinoculated in fresh broth was similar to that of bacteria from fresh culture or collected after the persister assay (Supplementary Figure 3), suggesting a reversion to a normal phenotype once the antibiotic pressure has been removed.

FACS Analysis of Intracellular Persisters

The S-LP (RN4220) and S-HP (1214) isolates expressing GFP under the control of a tetracycline-inducible promoter behaved intracellularly as clinical isolates, with C_s at values close to their MIC, and a lower (less negative) E_{max} of moxifloxacin when infected with S-HP than L-HP (-1.1 vs. $-2.3 \log_{10}$ CFU; Figure 4F). FACS analysis shows that propidium-negative (viable) bacteria collected from both strains after 24 h exposure to moxifloxacin at high concentrations remained highly fluorescent, indicating that they did not divide intracellularly (Figure 4G).

DISCUSSION

This study examines, in clinical *S. aureus* isolates exposed to the fluoroquinolone moxifloxacin, the correlation between (i) their high- vs. low-persister character (as determined by the relative abundance of persisters isolated from stationary cultures exposed to moxifloxacin), (ii) evolution to resistance (by spontaneous mutations upon exposure to subtherapeutic concentrations of the same antibiotic), and (iii) intracellular survival in permissive host cells in spite of exposure of these infected cells to high moxifloxacin concentrations. Isolates were from two geographically distinct origin (Vietnam and Belgium) to gain in clinically pertinent information and value. Conversely, we restricted this first analysis to a single antibiotic, taken as pharmacological tool, to minimize drug-related differences when addressing different biological and potentially disease-related properties of persisters. Moxifloxacin was selected as a highly bactericidal antibiotic, fulfilling

therefore the condition set for evidencing the occurrence of persisters (Balaban et al., 2019). In our set-up, it also offered the advantage of being clinically approved for the treatment of staphylococcal infections, while showing a balanced proportion of susceptible and resistant clinical isolates [a condition that becomes increasingly difficult to meet with other antibiotics when dealing with hospitalized patients, as exemplified here by our previous analysis of the resistance patterns of the Vietnamese collection (Nguyen et al., 2020)]. Three key observations are presented, which collectively may help in understanding how persistence to antibiotics may eventually account for clinical failures in spite of well-conducted antibiotic treatments.

First, the study shows a correlation between the ability of strains to give rise to persisters (quantitated here by their relative persister fraction) and their level of resistance to moxifloxacin. We indeed found that moxifloxacin-resistant isolates have all a high-persister character, while this character is more variable among susceptible isolates. This observation, as far as we know, has never been made using a large collection of clinical isolates. Antibiotic persistence and resistance are considered as distinct phenotypes leading independently to therapeutic failure (Fisher et al., 2017; Balaban et al., 2019). We show here, for moxifloxacin, that these two phenotypes are actually closely related and that resistance develops more rapidly in isolates with a high- rather than a low- relative persister fraction when exposed to subinhibitory concentrations of this antibiotic. Studies with laboratory strains of *E. coli* have demonstrated that persistence promotes evolution to resistance by increasing survival and mutation rates (Barrett et al., 2019; Windels et al., 2019). We extend this conclusion to *S. aureus* and clinical isolates. In this context, we add the observation that the expression of *recA*, the derepressor of the SOS response, is not induced in resistant isolates, but well in *in vitro* selected mutants, suggesting that the SOS response is involved in resistance acquisition but is no longer necessary once resistance has been acquired (Torres-Barcelo et al., 2015; Machuca et al., 2017). As expected, the main mechanism of resistance selected consisted in mutations in fluoroquinolones targets, with ParC and GyrA being most frequently affected, as in most other studies

(Schmitz et al., 1998; Roychoudhury et al., 2001; Kwak et al., 2013; Nakaminami et al., 2014).

In this context, a second key observation is that moxifloxacin induces the expression of genes encoding fluoroquinolone targets, especially in isolates with a high-persister character. This occurred independently of a decrease in susceptibility as well as its actual level, suggesting that this overexpression is associated with acquisition of a persister phenotype rather than of resistance. This is coherent with our recent observation that *gyrA* and *gyrB* are overexpressed in intracellular *S. aureus* persisters as part of a global response to the stress imposed by antibiotics, including by other classes than fluoroquinolones (Peyrusson et al., 2020).

A third major finding of this work is that isolates yielding a large proportion of persisters in stationary phase culture (broth) and phagocytized by THP-1 monocytes gave rise to larger residual inocula that remained unaffected by moxifloxacin intracellularly (less negative E_{\max} , which refers to the decrease in CFU's for an infinitely large extracellular concentrations, therefore denoting the drug maximal activity) compared to isolates categorized as showing a low-persister character in the same broth assay. We recently demonstrated for a laboratory strain of *S. aureus* that bacteria surviving to antibiotics intracellularly are persisters, showing a non-dividing phenotype readily reversible as soon as the antibiotic pressure is relieved (Peyrusson et al., 2020). We show here that this conclusion can be applied to clinical isolates as well, which opens a number of medically important avenues to explore. We also bring a rationale as why the intracellular maximal relative efficacy of a given antibiotic may vary among clinical isolates (Lemaire et al., 2011), if considering their potential differences in relative persister fraction. Thus, a high-persister character could stand as an important cause of ineffective elimination of intracellular *S. aureus*.

We also need to comment on the possible relationship between the presence of persisters among clinical isolates of *S. aureus* and persistence and/or recurrence of staphylococcal infections. Persisters are actually observed for all isolates from both the Vietnamese and Belgian collection (though in variable proportions), representing 0.001–1% of the population in stationary cultures if defined as cultivable organisms that remain unaffected by moxifloxacin at 100× its MIC. Conversely the residual intracellular inocula observed in cells exposed to similarly large concentrations of moxifloxacin represent from 1 to 10% of the post-phagocytosis inoculum, with most if not all being non-dividing, and therefore fulfilling in both respects the definition of persisters. This suggests a drastic enrichment in persisters when isolates are exposed to antibiotics in the intracellular milieu. If fractional, this effect of the antibiotics is anticipated to give rise to larger intracellular populations of persisters if infection is made with isolates that are producing more persisters in stationary phase cultures (and were categorized as having a high persister character). This is exactly what we found when comparing the sizes of the surviving intracellular pools (denoted by the differences in E_{\max} values), since these survivors were more numerous if the experiment was performed with isolates with

high relative persister fraction rather than with those with low relative persister fraction, even though the initial post-phagocytosis inocula were not different. This fully supports and strengthens the concept that one of the factors leading to intracellular survival in spite of exposure of the host cells to high antibiotic concentrations is the persister character of the isolate. Persisters are often considered as pre-existing to antibiotic exposure and be related to an ill-adapted immune host response or to the presence of growth-arrested bacteria (Zhang, 2014; Fisher et al., 2017). But we see here that the proportion of persisters is increased when the antibiotic exposure has taken place intracellularly. This, therefore, also suggests that the mechanisms responsible for the emergence of the persister phenotype involves more than pre-existing populations. It also reinforces our suggestion that persisters surviving antibiotic treatment intracellularly may contribute to relapses by recovering their capacity to divide and escape from the host cells once antibiotic therapy is discontinued (Fisher et al., 2017), since we see here that intracellular niches may result in considerable enrichment of the initial inocula with what are probably bacteria untreatable by currently available antibiotics. Resistance, however, remains important since moxifloxacin, at an extracellular concentration corresponding to its human C_{\max} (4 mg/L), had only a bacteriostatic effect on intracellular R-HP *S. aureus* isolates.

This study has limitations. First, we examined only a single antibiotic (moxifloxacin), which, however, was not only by design, as explained earlier, but also for feasibility reasons, having made the choice to work not with a limited number of strains but with a large collection of isolates to obtain conclusions that could be generalized to real-life samples and be potentially applicable to the clinic. Other fluoroquinolones could have been added to the study, but active and widely used anti-staphylococcal fluoroquinolones are few amongst the approved members of this class of antibiotics. Conversely, other bactericidal antibiotics from different pharmacological and/or chemical classes will need to be included in future investigations to reach truly generalizable conclusions. Of interest in this context, we recently observed that intracellular persisters can not only be selected by exposure to antibiotics belonging to different classes but also show unresponsiveness to multiple antibiotic classes, which can be explained by an activation of the global stress responses (Peyrusson et al., 2020). A second limitation stems from the lack of comprehensive molecular studies of the mechanisms underlying the differences in persister fractions among clinical isolates, which would require in-depth transcriptomic and/or genomic analyses, requiring, for critical cases, further documentation at the protein and function levels. Both limitations call for additional studies, but represent major undertakings by themselves.

At this stage, however, and in the limits of our current investigations, our work already opens the perspective that determining antibiotic persistence may help defining the risk of developing resistance and of maintaining a pool of dormant bacteria surviving intracellularly that may reactivate the infection at the end of the treatment. Screening for the persister character of clinical isolates, and establishing the predictive value and

specificity of the results of such assay, could become an important contribution to the improvement of antibiotic treatments in the future. This would require, however, the development of routine-usable methodologies, a first example of which could be the recently published TDtest, which used a modified disk-diffusion assay in order to detect antibiotic tolerance (Gefen et al., 2017).

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The study did not involve directly patients, but clinical isolates that were collected prospectively in Vietnam with the approval of the ethical committee from the Hanoi University of Pharmacy (1102/QD-DHN). The Belgian isolates were randomly selected in the collection of the Belgian reference center for Staphylococci, which does not need ethical approval.

AUTHOR CONTRIBUTIONS

TKN performed the experiments and analyzed the data. FP performed FACS analyses. MD and NHP provided clinical isolates and characterized them. HAN and PMT provided advice and PMT reanalyzed the data in details. FVB supervised the

study and wrote the Manuscript. All the authors made useful comments during the writing of the Manuscript and approved the submitted version.

FUNDING

This work was supported by the Belgian *Fonds de la Recherche Scientifique* FRS-FNRS (grants T.0189.16 and J.0018.17).

ACKNOWLEDGMENTS

TKN was successively junior research fellow from the *Fonds de coopération* of the *Université catholique de Louvain* and from the *Académie de Recherche et d'Enseignement Supérieur* (ARES). FP was recipient of a Ph.D. fellowship from the *Université catholique de Louvain*. FVB is Research Director from the Belgian *Fonds de la Recherche Scientifique* (FRS-FNRS). MD, NHP, and HAN acted within their normal academic, clinical or laboratory activities. PMT is emeritus and was unpaid. The authors are grateful to V. Mohymont and V. Yfantis for expert technical assistance.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2020.587364/full#supplementary-material>

REFERENCES

- Avrain, L., Garvey, M., Mesaros, N., Glupczynski, Y., Mingeot-Leclercq, M. P., Piddock, L. J. V., et al. (2007). Selection of quinolone resistance in *Streptococcus pneumoniae* exposed in vitro to subinhibitory drug concentrations. *J. Antimicrob. Chemother.* 60, 965–972. doi: 10.1093/jac/dkm292
- Balaban, N. Q., Helaine, S., Lewis, K., Ackermann, M., Aldridge, B., Andersson, D. I., et al. (2019). Definitions and guidelines for research on antibiotic persistence. *Nat. Rev. Microbiol.* 17, 441–448.
- Barcia-Macay, M., Seral, C., Mingeot-Leclercq, M. P., Tulkens, P. M., and Van Bambeke, F. (2006). Pharmacodynamic evaluation of the intracellular activities of antibiotics against *Staphylococcus aureus* in a model of THP-1 macrophages. *Antimicrob. Agents Chemother.* 50, 841–851. doi: 10.1128/aac.50.3.841-851.2006
- Barrett, T. C., Mok, W. W. K., Murawski, A. M., and Brynildsen, M. P. (2019). Enhanced antibiotic resistance development from fluoroquinolone persisters after a single exposure to antibiotic. *Nat. Commun.* 10:1177.
- Bateman, B. T., Donegan, N. P., Jarry, T. M., Palma, M., and Cheung, A. L. (2001). Evaluation of a tetracycline-inducible promoter in *Staphylococcus aureus* in vitro and in vivo and its application in demonstrating the role of sigB in microcolony formation. *Infect. Immun.* 69, 7851–7857. doi: 10.1128/iai.69.12.7851-7857.2001
- Bhagwat, S. S., Mundkur, L. A., Gupta, S. V., Patel, M. V., and Khorakiwala, H. F. (2006). The anti-methicillin-resistant *Staphylococcus aureus* quinolone WCK 771 has potent activity against sequentially selected mutants, has a narrow mutant selection window against quinolone-resistant *Staphylococcus aureus*, and preferentially targets DNA gyrase. *Antimicrob. Agents Chemother.* 50, 3568–3579. doi: 10.1128/aac.00641-06
- Boudjemaa, R., Steenkest, K., Jacqueline, C., Briandet, R., Caillon, J., Boutoille, D., et al. (2018). Live intramacrophagic *Staphylococcus aureus* as a potential cause of antibiotic therapy failure: observations in an in vivo mouse model of prosthetic vascular material infections. *J. Antimicrob. Chemother.* 73, 2418–2421. doi: 10.1093/jac/dky205
- Buyck, J. M., Lemaire, S., Seral, C., Anantharajah, A., Peyrusson, F., Tulkens, P. M., et al. (2016). In Vitro Models for the Study of the Intracellular Activity of Antibiotics. *Methods Mol. Biol.* 1333, 147–157. doi: 10.1007/978-1-4939-2854-5_13
- Cirz, R. T., Jones, M. B., Gingles, N. A., Minogue, T. D., Jarrahi, B., Peterson, S. N., et al. (2007). Complete and SOS-mediated response of *Staphylococcus aureus* to the antibiotic ciprofloxacin. *J. Bacteriol.* 189, 531–539. doi: 10.1128/jb.01464-06
- Cohen, N. R., Lobritz, M. A., and Collins, J. J. (2013). Microbial persistence and the road to drug resistance. *Cell Host. Microbe.* 13, 632–642. doi: 10.1016/j.chom.2013.05.009
- Conlon, B. P. (2014). *Staphylococcus aureus* chronic and relapsing infections: Evidence of a role for persister cells: An investigation of persister cells, their formation and their role in *S. aureus* disease. *Bioessays* 36, 991–996. doi: 10.1002/bies.201400080
- De Groote, V. N., Verstraeten, N., Fauvart, M., Kint, C. I., Verbeeck, A. M., Beullens, S., et al. (2009). Novel persistence genes in *Pseudomonas aeruginosa* identified by high-throughput screening. *FEMS Microbiol. Lett.* 297, 73–79.
- Dorr, T., Lewis, K., and Vulic, M. (2009). SOS response induces persistence to fluoroquinolones in *Escherichia coli*. *PLoS Genet.* 5:e1000760. doi: 10.1371/journal.pgen.1000760
- Dubourg, G., Abat, C., and Raoult, D. (2017). Why new antibiotics are not obviously useful now. *Int. J. Antimicrob. Agents* 49, 549–553. doi: 10.1016/j.ijantimicag.2016.11.015
- Fisher, R. A., Gollan, B., and Helaine, S. (2017). Persistent bacterial infections and persister cells. *Nat. Rev. Microbiol.* 15, 453–464. doi: 10.1038/nrmicro.2017.42

- Garzoni, C., and Kelley, W. L. (2009). *Staphylococcus aureus*: new evidence for intracellular persistence. *Trends Microbiol.* 17, 59–65. doi: 10.1016/j.tim.2008.11.005
- Gefen, O., Chekol, B., Strahilevitz, J., and Balaban, N. Q. (2017). TDtest: easy detection of bacterial tolerance and persistence in clinical isolates by a modified disk-diffusion assay. *Sci. Rep.* 7:41284.
- Heffernan, A. J., Sime, F. B., Lipman, J., and Roberts, J. A. (2018). Individualising Therapy to Minimize Bacterial Multidrug Resistance. *Drugs* 78, 621–641. doi: 10.1007/s40265-018-0891-9
- Jung, S. H., Ryu, C. M., and Kim, J. S. (2019). Bacterial persistence: Fundamentals and clinical importance. *J. Microbiol.* 57, 829–835. doi: 10.1007/s12275-019-9218-0
- Krut, O., Sommer, H., and Kronke, M. (2004). Antibiotic-induced persistence of cytotoxic *Staphylococcus aureus* in non-phagocytic cells. *J. Antimicrob. Chemother.* 53, 167–173. doi: 10.1093/jac/dkh076
- Kwak, Y. G., Truong-Bolduc, Q. C., Bin Kim, H., Song, K. H., Kim, E. S., and Hooper, D. C. (2013). Association of *norB* overexpression and fluoroquinolone resistance in clinical isolates of *Staphylococcus aureus* from Korea. *J. Antimicrob. Chemother.* 68, 2766–2772. doi: 10.1093/jac/dkt286
- Lemaire, S., Glupczynski, Y., Duval, V., Joris, B., Tulkens, P. M., and Van Bambeke, F. (2009). Activities of ceftobiprole and other cephalosporins against extracellular and intracellular (THP-1 macrophages and keratinocytes) forms of methicillin-susceptible and methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 53, 2289–2297. doi: 10.1128/aac.01135-08
- Lemaire, S., Kosowska-Shick, K., Appelbaum, P. C., Glupczynski, Y., Van Bambeke, F., and Tulkens, P. M. (2011). Activity of moxifloxacin against intracellular community-acquired methicillin-resistant *Staphylococcus aureus*: comparison with clindamycin, linezolid and co-trimoxazole and attempt at defining an intracellular susceptibility breakpoint. *J. Antimicrob. Chemother.* 66, 596–607. doi: 10.1093/jac/dkq478
- Machuca, J., Recacha, E., Biales, A., Diaz-de-Alba, P., Blazquez, J., Pascual, A., et al. (2017). Cellular Response to Ciprofloxacin in Low-Level Quinolone-Resistant *Escherichia coli*. *Front. Microbiol.* 8:1370. doi: 10.3389/fmicb.2017.01370
- Martinez, J. L., Baquero, F., and Andersson, D. I. (2007). Predicting antibiotic resistance. *Nat. Rev. Microbiol.* 5, 958–965.
- Maslowska, K. H., Makiela-Dzbenka, K., and Fijalkowska, I. J. (2019). The SOS system: A complex and tightly regulated response to DNA damage. *Environ. Mol. Mutagen.* 60, 368–384. doi: 10.1002/em.22267
- Morrow, B. J., Abbanat, D., Baum, E. Z., Crespo-Carbone, S. M., Davies, T. A., He, W., et al. (2011). Antistaphylococcal activities of the new fluoroquinolone JNJ-Q2. *Antimicrob. Agents Chemother.* 55, 5512–5521. doi: 10.1128/aac.00470-11
- Nakaminami, H., Sato-Nakaminami, K., and Noguchi, N. (2014). A novel *GyrB* mutation in methicillin-resistant *Staphylococcus aureus* (MRSA) confers a high level of resistance to third-generation quinolones. *Int. J. Antimicrob. Agents* 43, 478–479. doi: 10.1016/j.ijantimicag.2014.02.002
- Nguyen, T. K., Argudin, M. A., Deplano, A., Pham, N. H., Nguyen, H. A., Tulkens, P. M., et al. (2020). Antibiotic resistance, biofilm formation, and intracellular survival as possible determinants of persistent or recurrent infections by *Staphylococcus aureus* in a Vietnamese tertiary hospital. Focus on bacterial response to moxifloxacin. *Microb. Drug Resist.* 26, 537–544. doi: 10.1089/mdr.2019.0282
- Noguchi, N., Okihara, T., Namiki, Y., Kumaki, Y., Yamanaka, Y., Koyama, M., et al. (2005). Susceptibility and resistance genes to fluoroquinolones in methicillin-resistant *Staphylococcus aureus* isolated in 2002. *Int. J. Antimicrob. Agents* 25, 374–379. doi: 10.1016/j.ijantimicag.2004.11.016
- Ortiz-Padilla, M., Diaz-Diaz, S., Machuca, J., Tejada-Gonzalez, A., Recacha, E., Docobo-Perez, F., et al. (2020). Role of low-level quinolone resistance in generating tolerance in *Escherichia coli* under therapeutic concentrations of ciprofloxacin. *J. Antimicrob. Chemother.* 75, 2124–2132.
- Peyrusson, F., Nguyen, T. K., Varet, H., Legendre, R., Sismeiro, O., Coppée, J. Y., et al. (2020). Intracellular *Staphylococcus aureus* persists upon antibiotic exposure. *Nat. Commun.* 11:2200.
- Peyrusson, F., Van Wessem, A., Dieppois, G., Van Bambeke, F., and Tulkens, P. M. (2019). Cellular pharmacokinetics and intracellular activity of the bacterial fatty acid synthesis inhibitor, afabacin desphosphono against different resistance phenotypes of *Staphylococcus aureus* in models of cultured phagocytic cells. *Int. J. Antimicrob. Agents* 55:105848. doi: 10.1016/j.ijantimicag.2019.11.005
- Qin, T. T., Kang, H. Q., Ma, P., Li, P. P., Huang, L. Y., and Gu, B. (2015). SOS response and its regulation on the fluoroquinolone resistance. *Ann. Transl. Med.* 3:358.
- Rhoads Kraemer, G., and Iandolo, J. J. (1990). High-frequency transformation of *Staphylococcus aureus* by electroporation. *Curr. Microbiol.* 21, 373–376. doi: 10.1007/bf02199440
- Roychoudhury, S., Twinem, T. L., Makin, K. M., Nienaber, M. A., Li, C., Morris, T. W., et al. (2001). *Staphylococcus aureus* mutants isolated via exposure to nonfluorinated quinolones: detection of known and unique mutations. *Antimicrob. Agents Chemother.* 45, 3422–3426. doi: 10.1128/aac.45.12.3422-3426.2001
- Schmitz, F. J., Hofmann, B., Hansen, B., Scheuring, S., Luckefahr, M., Klootwijk, M., et al. (1998). Relationship between ciprofloxacin, ofloxacin, levofloxacin, sparfloxacin and moxifloxacin (BAY 12-8039) MICs and mutations in *griA*, *griB*, *gyrA* and *gyrB* in 116 unrelated clinical isolates of *Staphylococcus aureus*. *J. Antimicrob. Chemother.* 41, 481–484. doi: 10.1093/jac/41.4.481
- Torres-Barcelo, C., Kojadinovic, M., Moxon, R., and MacLean, R. C. (2015). The SOS response increases bacterial fitness, but not evolvability, under a sublethal dose of antibiotic. *Proc. Biol. Sci.* 282:20150885. doi: 10.1098/rspb.2015.0885
- Tuschscherr, L., Löffler, B., and Proctor, R. A. (2020). Persistence of *Staphylococcus aureus*: Multiple Metabolic Pathways Impact the Expression of Virulence Factors in Small-Colony Variants (SCVs). *Front. Microbiol.* 11:1028. doi: 10.3389/fmicb.2020.01028
- Van den Bergh, B., Fauvart, M., and Michiels, J. (2017). Formation, physiology, ecology, evolution and clinical importance of bacterial persisters. *FEMS Microbiol. Rev.* 41, 219–251. doi: 10.1093/femsre/fux001
- Windels, E. M., Michiels, J. E., Fauvart, M., Wenseleers, T., Van den Bergh, B., and Michiels, J. (2019). Bacterial persistence promotes the evolution of antibiotic resistance by increasing survival and mutation rates. *ISME J.* 13, 1239–1251. doi: 10.1038/s41396-019-0344-9
- Zhang, Y. (2014). Persisters, persistent infections and the Yin-Yang model. *Emerg. Microbes. Infect.* 3:e3.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Nguyen, Peyrusson, Dodémont, Pham, Nguyen, Tulkens and Van Bambeke. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Supplementary Material

Table S1: characteristics of the strains used in this study

S. aureus ATCC 25923 and RN 4220 are reference strains from the American Tissue Culture Collection (ATCC), Manassas, VA. Thirty-six clinical isolates were collected at the Bach Mai Hospital (Hanoi, Vietnam) from patients with microbiological and clinical evidence of persisting and/or recurrent infection (defined as (i) an infection that did not resolve after 5 days of treatment by antibiotics to which the initial isolates were reported as susceptible or (ii) an infection that reactivated a few days after the administration of an antibiotic to which the initial isolate was susceptible but had been discontinued based on clinical evidence of improvement of the patient's conditions)(Nguyen et al., 2020). Their Agr group, typing (based on *spa* [Staphylococcus protein A] type and Pulsed Field Gel Electrophoresis), and susceptibility to 26 anti-staphylococcal antibiotics (covering the main pharmacological classes) have been reported previously (Nguyen et al., 2020). Eighteen randomly-selected isolates from the collection of the Belgian National Reference Center for Staphylococci were included as complement.

Origin	Strain identification	MSSA/MRSA ^a	Moxifloxacin MIC (mg/L) ^b	Moxifloxacin relative persister fraction (RPF) ^c	Assigned phenotypic group
References	ATCC 25923	MSSA	0.032	1.0	Susceptible with low– relative persister fraction (S-LP)
	RN 4220	MSSA	0.064	8.0	
Bach Mai hospital, Hanoi, Vietnam	69474	MRSA	0.064	0.2	Susceptible with low– relative persister fraction (S-LP)
	30549	MSSA	0.032	0.8	
	3446	MSSA	0.064	0.8	
	69687	MRSA	0.25	1.0	
	30337	MRSA	0.064	1.2	
	69493	MRSA	0.064	1.2	
	22714	MRSA	0.064	1.3	
	69613	MSSA	0.064	1.4	
	69486	MRSA	0.064	1.7	
	17883	MSSA	0.032	2.2	
	30461	MSSA	0.064	2.4	
	19255	MSSA	0.064	2.5	
	69700	MRSA	0.064	3.9	
	34613	MRSA	0.032	6.2	

	69544	MRSA	0.032	7.8	Susceptible with high– relative persister fraction (H-LP)
	34724	MRSA	0.064	8.6	
	36271	MRSA	0.032	9.0	
	36606	MSSA	0.032	12.9	
	14305	MSSA	0.032	24.8	
	69783	MSSA	0.032	37.2	
	69505	MRSA	0.032	45.5	
	69867	MRSA	0.064	74.8	
	30566	MSSA	2	60.7	Resistant with high– relative persister fraction (R-HP)
	30342	BORSA or MODSA	2	79.8	
	69519	MRSA	1	91.6	
	20975	MSSA	2	92.7	
	30609	MRSA	2	100.8	
	25619	MRSA	2	111.3	
	69781	MSSA	2	138.2	
	36253	MRSA	2	176.0	
	30462	MSSA	2	176.9	
	34791	BORSA or MODSA	2	196.8	
	35994	MSSA	2	228.1	
	69915	MRSA	1	273.1	
	13890	MSSA	1	289.6	
	34427	MSSA	2	411.8	
Reference Center for Staphylococci, Brussels, Belgium	2013S467	MRSA	0.064	0.4	Susceptible with low– relative persister fraction (S-LP)
	2014S449	MSSA	0.032	0.8	
	2017S228	MRSA	0.064	1.2	
	1078	MSSA	0.064	1.5	
	2016S354	MRSA	0.064	1.7	
	1077	MSSA	0.064	1.7	
	2014S092	MRSA	0.032	2.2	
	2014S118	MRSA	0.064	2.4	
	238	MSSA	0.064	2.9	
	2015S580	MSSA	0.064	3.0	
	1159	MSSA	0.064	3.0	
	196	MRSA	0.032	6.7	

	1214	MSSA	0.032	26.7	Susceptible with high– relative persister fraction (S-HP)
	2015S418	MRSA	1	53.6	Resistant with high– relative persister fraction (R-HP)
	2016S605	MRSA	2	154.8	
	2013S092	MRSA	1	221.7	
	2017S351	MRSA	2	250.6	
	2017S017	MRSA	2	301.4	

^a determined based on PCR of *mecA* and *mecC* genes and MIC of cefoxitine

^b categorization as susceptible (MIC \leq 0.25 mg/L) or resistant (MIC $>$ 0.25 mg/L) was made according to the 2019 interpretive criteria of the European Committee for Antibiotic Susceptibility Testing (<https://www.eucast.org>).

^c ratio between the percentage of persisters after 5 h incubation with moxifloxacin at 100 x MIC for the clinical isolate and for ATCC25923; values are mean of 3 independent experiments performed in triplicates

Table S2: primers used in this study RT-PCR and PCR experiments

Gene	Primer	Primer sequence (5'→3')	Purpose	Product size (bp)	Reference
<i>gmk</i> [§]	gmk-F	TCAGGACCATCTGGAGTAGGTAAAG	qRT-PCR	108	(Truong-Bolduc et al., 2012)
	gmk-R	TTCACGCATTTGACGTGTTG	qRT-PCR		
<i>recA</i>	recA-F	GTAGCGCTTCACGCTATTGCT	qRT-PCR	not disclosed	(Cirz et al., 2007)
	recA-R	TTCAAGACCTTGTTACCATGATC	qRT-PCR		
<i>lexA</i>	lexA-F	GAAACGATTCATGTGCCAGTTATT	qRT-PCR	not disclosed	(Cirz et al., 2007)
	lexA-R	GTCGCTATTATGTGTCGATGTAA	qRT-PCR		
<i>uvrB</i>	uvrB-F	AATATTCCCAGCCTCTAAAGAAGAA	qRT-PCR	not disclosed	(Anderson et al., 2006)
	uvrB-R	CTCATCTCGTAATTCTTTCAATCGT	qRT-PCR		
<i>umuC</i>	umuC-F	TGCGAGTGTTTCTTGTATTG	qRT-PCR	not disclosed	(Schroder et al., 2013)
	umuC-R	CCCTGTCTTGATGCCTAA	qRT-PCR		
<i>gyrA</i>	RT-gyrA-F	TCGTGAAGGTGACGAAGTTG	qRT-PCR	101	This study ^a
	RT-gyrA-R	CTGGCGTACGTTTACCATAAC	qRT-PCR		
<i>gyrB</i>	RT-gyrB-F	GGTGCTGGGCAAATACAAGT	qRT-PCR	107	
	RT-gyrB-R	TCCCACACTAAATGGTGCAA	qRT-PCR		
<i>parC</i>	RT-parC-F	GCGTGACGCTAAAGAAAACC	qRT-PCR	121	
	RT-parC-R	TTCACCTTCAAGCGCAACTA	qRT-PCR		
<i>parE</i>	RT-parE-F	GACGTGGTATGCCAACAGGT	qRT-PCR	124	
	RT-parE-R	CCGTGAAGACCACCTGAAGT	qRT-PCR		
<i>norA</i>	RT-norA-F	GACATTTACCAAGCCATCAA	qRT-PCR	102	
	RT-norA-R	TGCCATAAATCCACCAATCC	qRT-PCR		
<i>norB</i>	RT-norB-F	GCTACACCATCAACAGATACAGCAA	qRT-PCR	117	
	RT-norB-R	ACTCAATGCGACGCCAAA	qRT-PCR		
<i>norC</i>	RT-norC-F	TGGGTTGGAGATGGATTTTC	qRT-PCR	130	
	RT-norC-R	ACAATTAGCCCTGCAACGTC	qRT-PCR		

<i>gyrA</i>	Seq-gyrA-F	AATGAACAAGGTATGACACC	PCR	223	(Schmitz et al., 2002)
	Seq-gyrA-R	TACGCGCTTCAGTATAACGC	PCR		
<i>gyrB</i>	Seq-gyrB-F	CAGCGTTAGATGTAGCAAGC	PCR	250	
	Seq-gyrB-R	CCGATTCCTGTACCAAATGC	PCR		
<i>parC</i>	Seq-parC-F	ACTTGAAGATGTTTTAGGTGAT	PCR	459	
	Seq-parC-R	TTAGGAAATCTTGATGGCAA	PCR		
<i>parE</i>	Seq-parE-F	CGATTAAAGCACAACAAGCAAG	PCR	375	
	Seq-parE-R	CATCAGTCATAATAATTACTC	PCR		

^a designed using Primer3 Software (<http://bioinfo.ut.ee/primer3-0.4.0/>), using the published sequences of *griA* (also referred to as *parC*) and *griB* also referred to as *parE* (Yamagishi et al., 1996) and *gyrA* and *gyrB* (Ito et al., 1994) of *S. aureus* RN4220.

Table S3: Mutations in QRDR domains of selected isolates^a

Phenotype	Isolates	MIC	<i>gyrA</i>	<i>gyrB</i>	<i>parC</i>	<i>parE</i>
reference	RN4220	0.064	-	-	-	-
S-LP isolates	30337 (i)	0.064	-	-	-	-
	30337* (f) – 32 days	2		<i>Leu433Ile^b</i>	<i>Ser80Phe^{b,d}</i>	<i>Glu422Asp^c</i>
	69474 (i)	0.064	-	-	-	Glu422Asp
	69474* (f) – 35 days	2		<i>Ile454Val^b</i>	<i>Ser80Ala^b</i>	<i>Glu422Asp</i>
	69687 (i)	0.25	-	-	His103Tyr ^{b,d}	Glu422Asp
	69687* (f) -18 days	2	<i>Ser84Leu^{b,d}</i>		<i>His103Tyr</i>	<i>Glu422Asp</i>
S-HP isolates	69505 (i)	0.032	-	-	-	Glu422Asp
	69505* (f) – 15 days	2	<i>Ser84Leu</i>	-	-	<i>Glu422Asp</i>
	69783 (i)	0.032	-	-	-	-
	69783* (f)- 16 days	2	<i>Ser84Leu</i>	-	-	<i>Glu422Asp</i>
	69867 (i)	0.064	-	-	-	Glu422Asp
	69867* (f) -19 days	2	<i>Ser84Leu</i>	-	-	<i>Glu422Asp</i>
R-HP isolates	30642	2	Ser84Leu		Ser80Phe	-
	69781	2	Ser84Leu	-	Ser80Phe	-
	30566	2	Ser84Leu	-	Ser80Phe	-
	34427	2	Ser84Leu	-	Ser80Phe	-
	13890	1	Ser84Leu	-	Ser80Phe	-
	35994	2	Ser84Leu	-	Ser80Phe	-

^a only non-silent mutations are shown^b base changes leading to amino acid substitutions: Ser84Leu: TCA→TTA; Leu433Ile: TTA→ATA; Ile454Val: ATT→GTT; Ser80Phe: TCC→TTC; Ser80Ala: TCC→GCC; His103Tyr: CAT→TAT; Glu422Asp: GAA→GAT^c mutations described as not associated with resistance (Schmitz et al., 1998)^d mutations described as associated with resistance (Kwak et al., 2013; Morrow et al., 2011; Roychoudhury et al., 2001).


* in-vitro selected mutants (f) from the corresponding parental isolates (i), after exposure to ½ MIC of moxifloxacin until reaching an MIC of 2 mg/L. The number of days of subcultures needed to reach this MIC is indicated after the name of the isolate.

Table S4: Key pharmacodynamic parameters of the concentration-effects experiments shown in Figure 4: values for (i) each individual strain (single experiments with triplicate determinations of cfu's for each extracellular concentration (30 in total) and fitting of one Hill-Langmuir function (with slope factor set to 1) and (ii) all strains within the same phenotypic group analyzed together (240, 150, or 120 data points, respectively) and used for fitting a single Hill-Langmuir function (with slope factor set to 1) *. All data are shown as the mean value and the corresponding 95% confidence interval (asymmetrical range of values containing the true mean with 95% of certainty)

S-LP^a			S-HP^a			R-HP^a		
isolate	IC₅₀^b	E_{max}^c	isolate	IC₅₀^b	E_{max}^c	isolate	IC₅₀^b	E_{max}^c
2014S118	1.11 0.83 to 1.48	-2.089 -2.39 to -1.80	1214	0.862 0.58 to 1.31	-1.12 -1.45 to -0.78	2015S418	1.01 0.64 to 1.63	-1.46 -1.86 to -1.07
2016S354	1.44 0.85 to 2.55	-2.58 -3.10 to -2.04	69505	1.29 0.73 to 2.34	-1.54 -1.90 to -1.20	2017S017	0.73 0.48 to 1.10	-1.37 -1.73 to -1.03
2017S228	0.80 0.54 to 1.18	-1.78 -2.11 to -1.45	69783	1.92 1.14 to 3.37	-1.84 -2.09 to -1.24	20975	1.30 0.68 to 2.68	-1.59 -2.12 to -1.12
1078	1.00 0.620 to 1.66	-1.59 -2.03 to -1.19	69867	3.01 1.36 to 7.52	-1.61 -2.24 to -1.47	13890	2.68 1.76 to 4.14	-0.903 -1.19 to -0.630
30337	1.07 0.673 to 1.76	-2.18 -2.61 to -1.78	36606	1.36 0.83 to 2.30	-1.12 -1.47 to -0.780			
69474	1.76 0.93 to 3.58	-2.265 -2.81 to -1.78						
69687	2.11 1.45 to 3.09	-2.47 -2.85 to -2.11						
34467	0.833 0.55 to 1.28	-1.830 -2.22 to -1.46						
all^d	1.17 0.98 to 1.39	-2.10 -2.38 to -1.81	all^d	1.35 0.99 to 1.87	-1.45 -1.84 to -1.06	all^d	1.14 0.856 to 1.53	-1.33 -1.80 to -0.85

- * see the figure associated to this Table for residual QQ plots checking that the distributions of residuals are Gaussian for all sets of data.
- ^a Phenotype: S-LP: susceptible and low persister; S-HP: susceptible and high persister; R-HP: resistant and high persister (see main text for definition of susceptible/resistant and low persister/high persister phenotypes).
- ^b moxifloxacin extracellular concentration (multiples of the MIC for each strain) causing a change in cfu's half-way between E_{\min} and E_{\max} (defined hereunder)
- ^c E_{\max} : maximal relative efficacy (decrease of intracellular cfus' from the post-phagocytosis intracellular inoculum for an infinitely large extracellular moxifloxacin concentration; E_{\min} : minimal relative efficacy (increase of intracellular cfu's from post-phagocytosis intracellular inoculum for an infinitely low moxifloxacin extracellular concentration (both calculated using the Hill-Langmuir function (with slope factor set to 1) fitted to the data).
- ^d fit of all data from all strains in a given phenotype to a single Hill-Langmuir function (with slope factor set to 1). Comparison between models involving (i) different Hill-Langmuir functions (each fitted to the data of a single strain) vs (ii) a single function fitted to the data of all strains in each phenotype group, using the Akaike's information criteria (AIC) showed a probability of >99% for the single function model (differences in AIC: - 119.1, -55.6 and -56.49 for the S-LP, S-HP, and R-HP phenotypic groups, respectively when comparing it with the multiple functions model.

Residual QQ plots (for assessment of Gaussian distribution * for the least square regression made to fit Hill-Langmuir functions to the data of the concentration-effects experiments illustrated in Figure 4 and for which the values of the IC_{50} and E_{max} parameters are shown in Table S4). Left panels: plots for distinct Hill-Langmuir function fitted to the data of each individual strain (each symbol corresponds to one strain). Right panels: plots for a single Hill-Langmuir function fitted to the data of all strains with a similar phenotype.



* X axis: actual residual; Y axis: data points are very close to the line of identity, shown in red on the graph, consistent with a Gaussian distribution of the data used for the regression analysis.

Figure S1

Expression of the genes encoding proteins involved in SOS response (A) or fluoroquinolone targets (B) in selected isolates of *S. aureus* with susceptible low-persister (S-LP(i), red [30337, 69474, 69687]), susceptible high-persister (S-HP(i), green [69505, 69783, 69867]) phenotypes and in their resistant mutants selected in-vitro and having reached an MIC of 2 mg/L (S-LP(f) and S-HP(f)) after culture with subMIC concentrations of moxifloxacin (Figure 3). Isolates were incubated for 5 h with moxifloxacin at 100 x MIC in the conditions used for performing the persister assay (Figure 1). Each dot represents the mean of triplicates for one isolate, with horizontal lines showing the mean and SD. Data are expressed as $-\Delta\Delta CT$, with the value of ATCC 25923 at time 0 h used as a reference (horizontal dotted line). The thin horizontal line shows the expression level measured for ATCC 25923 at this time point. Statistical analysis: paired t test comparing S-LP(i) and S-HP(i) to S-LP(f) and S-HP(f), respectively: * : $p < 0.05$; **: $p < 0.01$





Figure S2

Expression of the genes encoding fluoroquinolone efflux pumps in selected isolates of *S. aureus* with susceptible low-persister (S-LP(i), red [30337, 69474, 69687]), susceptible high-persister (S-HP(i), green [69505, 69783, 69867]) or resistant high-persister (R-HP, blue [30462, 69781, 30566, 13890, 35994]) phenotypes as well as in mutants selected in-vitro from S-LP(i) and S-HP(i) isolates and having reached an MIC of 2 mg/L (S-LP(f) and S-HP(f)) after culture with subMIC concentrations of moxifloxacin (Figure 3). Isolates were incubated for 1h with moxifloxacin at 100xMIC in the conditions used for performing the persister assay (Figure 1). Similar results were obtained after 5 h of incubation. Each dot represents the mean of triplicates for one isolate, with horizontal lines showing the mean and SD. Data are expressed as $-\Delta\Delta CT$, with the value of ATCC 25923 at time 0 h used as a reference (horizontal dotted line). The thin horizontal line in panels B and C shows the expression level measured for ATCC 25923 at 1h. Statistical analysis: The figures at the top of the graphs shows the p value of a t-test comparing the group of isolates with ATCC 25923 at time 0h. p values lower than 0.05 are squared in red. Paired t test comparing S-LP(i) and S-HP(i) to S-LP(f) and S-HP(f) or ANOVA comparing parental isolates from different phenotypic subgroups revealed no statistical differences.



References to the supplementary material

- Anderson, K. L., Roberts, C., Disz, T., Vonstein, V., Hwang, K., Overbeek, R., Olson, P. D., Projan, S. J., and Dunman, P. M. (2006). Characterization of the *Staphylococcus aureus* heat shock, cold shock, stringent, and SOS responses and their effects on log-phase mRNA turnover. *J. Bacteriol.* 188, 6739-6756.
- Cirz, R. T., Jones, M. B., Gingles, N. A., Minogue, T. D., Jarrahi, B., Peterson, S. N., and Romesberg, F. E. (2007). Complete and SOS-mediated response of *Staphylococcus aureus* to the antibiotic ciprofloxacin. *J. Bacteriol.* 189, 531-539.
- Ito, H., Yoshida, H., Bogaki-Shonai, M., Niga, T., Hattori, H., and Nakamura, S. (1994). Quinolone resistance mutations in the DNA gyrase *gyrA* and *gyrB* genes of *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 38, 2014-2023.
- Kwak, Y. G., Truong-Bolduc, Q. C., Bin Kim, H., Song, K. H., Kim, E. S., and Hooper, D. C. (2013). Association of *norB* overexpression and fluoroquinolone resistance in clinical isolates of *Staphylococcus aureus* from Korea. *J. Antimicrob. Chemother.* 68, 2766-2772.
- Morrow, B. J., Abbanat, D., Baum, E. Z., Crespo-Carbone, S. M., Davies, T. A., He, W., Shang, W., Queenan, A. M., and Lynch, A. S. (2011). Antistaphylococcal activities of the new fluoroquinolone JNJ-Q2. *Antimicrob. Agents Chemother.* 55, 5512-5521.
- Nguyen, T. K., Argudin, M. A., Deplano, A., Pham, N. H., Nguyen, H. A., Tulkens, P. M., Dodemont, M., and Van Bambeke, F. (2020). Antibiotic resistance, biofilm formation, and intracellular survival as possible determinants of persistent or recurrent infections by *Staphylococcus aureus* in a Vietnamese tertiary hospital. Focus on bacterial response to moxifloxacin. *Microb. Drug Resist.* 26, 537-544.
- Roychoudhury, S., Twinem, T. L., Makin, K. M., Nienaber, M. A., Li, C., Morris, T. W., Ledoussal, B., and Catrenich, C. E. (2001). *Staphylococcus aureus* mutants isolated via exposure to nonfluorinated quinolones: detection of known and unique mutations. *Antimicrob. Agents Chemother.* 45, 3422-3426.
- Schmitz, F. J., Hofmann, B., Hansen, B., Scheuring, S., Luckefahr, M., Klootwijk, M., Verhoef, J., Fluit, A., Heinz, H. P., Kohrer, K., and Jones, M. E. (1998). Relationship between ciprofloxacin, ofloxacin, levofloxacin, sparfloxacin and moxifloxacin (BAY 12-8039) MICs and mutations in *grlA*, *grlB*, *gyrA* and *gyrB* in 116 unrelated clinical isolates of *Staphylococcus aureus*. *J. Antimicrob. Chemother.* 41, 481-484.
- Schmitz, F. J., Boos, M., Mayer, S., Kohrer, K., Scheuring, S., and Fluit, A. C. (2002). In vitro activities of novel des-fluoro(6) quinolone BMS-284756 against mutants of *Streptococcus pneumoniae*, *Streptococcus pyogenes*, and *Staphylococcus aureus* selected with different quinolones. *Antimicrob. Agents Chemother.* 46, 934-935.

- Schroder, W., Goerke, C., and Wolz, C. (2013). Opposing effects of aminocoumarins and fluoroquinolones on the SOS response and adaptability in *Staphylococcus aureus*. *J. Antimicrob. Chemother.* 68, 529-538.
- Truong-Bolduc, Q. C., Hsing, L. C., Villet, R., Bolduc, G. R., Estabrooks, Z., Taguezem, G. F., and Hooper, D. C. (2012). Reduced aeration affects the expression of the NorB efflux pump of *Staphylococcus aureus* by posttranslational modification of MgrA. *J. Bacteriol.* 194, 1823-1834.
- Yamagishi, J., Kojima, T., Oyamada, Y., Fujimoto, K., Hattori, H., Nakamura, S., and Inoue, M. (1996). Alterations in the DNA topoisomerase IV *grlA* gene responsible for quinolone resistance in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 40, 1157-1163.