# Poster A1-677

# Contrasting Effect of Acidic pH on the Extracellular and Intracellular Activities of Delafloxacin (DFX) vs Moxifloxacin (MXF) Towards S. aureus

# 50th ICAAC **Boston, MA, USA** September 12-15, 2010

#### ABSTRACT

contrast to currently marketed fluoroquinolones (FQ), DFX (1-(6-Amino-3,5-difluoro-pyridin-2-yl) -8-chloro-6-fluoro-7-(3-hydroxyazetidin-1-yl)-4-oxo-1,4-dihydro- quinoline-3-carboxylic acid) highly active on Gram-positive bacteria, is an anionic compound (pKa's = 0.80 and 5.49). In this context, we have examined the effect of acidic pH on its accumulation in *S. aureus* and in human THP-1 cells, in parallel with its activity against extra- and intracellular *S. aureus* (localized in phagolysosomes, pH 5-5.5). MXF (pKa's = 6.04 and 10.61) was used as comparator.

**Methods:** MICs measured in pH-adjusted Mueller-Hinton broth by microdilution. FQ accumulation in S. aureus ATCC 25923 and in uninfected THP-1 cells determined by microbiological assay. Intracellular activity assessed after 24 h of incubation of cells in media at pH 7.4 or 5.5 (JAC 2005, 55:897-904).

**Results:** The Table shows that DFX is generally more potent than MXF, as it shows activity at lower conc. both extracell. and intracell. At reduced pH, this difference is enhanced as acidification (i) increases about 10-fold DFX accumulation in both bacteria and cells, and (ii) decreases DFX MIC (5 dil.), resulting in a marked increase in intracell. activity (0.8-2.4 log10). Opposite effects are seen for MXF.

**Conclusions:** DFX displays a markedly improved potency and activity at acidic pH, probably related to its anionic character and ensuing increased accumulation in both bacteria and eucaryotic cells. This may confer an advantage to DFX for the treatment of staphylococcal infections in territories where pH is acidic.

Parameters studied	Moxifloxacin		Delafloxacin	
	pH 7.4	pH 5.5	pH 7.4	pH 5.5
Accumulation in	100.0 ± 4.1	57.8 ± 4.7	100.0 ± 9.7	1100.2 ± 180.7
S. aureus <sup>a</sup>				
MICs (mg/L)	0.03	0.125	0.0075	0.0002
Accumulation in	$59\pm05$	1.5 ± 0.0	1.7 ± 0.4	19.8 ± 4.7
THP-1 cells <sup>b</sup>	$3.0 \pm 0.3$			
Intracell. activity <sup>c</sup>				
at 0.1 mg/L	$0.7 \pm 0.2$	$0.5 \pm 0.0$	-0.5 ± 0.01	$-1.3 \pm 0.0$
at 1 mg/L	-1.5 ± 0.1	0.1 ± 0.1	$-0.6 \pm 0.0$	-3.0 ± 0.1
5				

<sup>a</sup> Antibiotic uptake within *S. aureus* (30 min.; % of control value [pH 7.4])

<sup>2</sup> Cellular to extracellular concentration ratio (30 min.)

<sup>c</sup>  $\Delta$ log cfu from time 0 (24 h)

#### BACKGROUND

Delafloxacin (DFX) is an investigational quinolone demonstrating potent intrinsic activity, especially against Grampositive bacteria (including MRSA isolates)<sup>1-2</sup>. From a chemical point of view, delafloxacin differs from other currently used quinolones by the absence of a protonable substituant, which confers a weak acid character to the molecule. This property further increases its potency in acidic environments. This could be advantageous for S. aureus infections, as this bacterium has ability to survive and thrive in mildly acidic compartments<sup>3-4</sup> (such as the skin, the vagina, the urinary tract, or the phagolysosomes of infected cells [pH 5-5.5]).

Structural formulae: The open arrows point to the acidic function of fluoroquinolones and the plain arrow, to the basic function of moxifloxacin that is protonated at physiological pH.





Sandrine Lemaire, Paul M. Tulkens, and Françoise Van Bambeke

Unité de Pharmacologie cellulaire et moléculaire, Louvain Drug Research Institute, Université catholique de Louvain, Brussels, Belgium

# **STUDY AIM**

To examine the effect of acidic pH: -on the accumulation of delafloxacin and moxifloxacin in procaryotic cells.

-on the accumulation of delafloxacin and moxifloxacin in eucaryotic cells.

-on the activity of delafloxacin and moxifloxacin towards extracellular and intracellular S. aureus.

#### METHODS

Bacteria, susceptibility testing, and extracellular activity: S. aureus strain ATCC 25923 was used for all experiments. MICs were determined in pH-adjusted Mueller Hinton Broth by microdilution. Extracellular activity was determined as previously described (4).

Cell line: Experiments were performed with THP-1 cells (ATCC TIB-202), a human myelomonocytic cell line displaying macrophage-like activity.

Determination of antibiotic accumulation in S. aureus and THP-1 cells: Antibiotic accumulation was measured after 30 min. incubation. The content in antibiotic was determined by microbiological assay (disc-plate). The cellular concentration of fluoroquinolones was expressed by reference to the total cell protein, as determined using the Folin-Ciocalteu/biuret method. Cellular accumulation in THP-1 cells was calculated using a conversion factor of 5 μL per mg of cell protein.

Cell infection and assessment of intracellular activity: Cell infection was performed exactly as described previously (4) with a starting inoculum of ~ 1 to 2 x  $10^{6}$  CFU/mg protein. Antibiotic activities was examined after 24 hours in the presence of fixed concentration of antibiotics in media at pH 7.4 or 5.5, and results were expressed as the change in the inoculum at 24 hours compared to time zero.

### REFERENCES

- 1) Ohshita et al, abstr. F-84, 38th ICAAC (1998)
- 2) Burak et al, P-1080, 19th ECCMID (2009)
- 3) Weinrick et al, J Bacteriol (2004), 186:8407-23
- 4) Barcia-Macay et al, Antimicrob Agents Chemother. (2006), 50:841-51

### ACKNOWLEDGEMENTS

ACKNOWLEDGEMENTS: We thank M.C. Cambier and C. Misson for technical assistance. S.L. and F.V.B. are Post-doctoral and Senior Research Associate of the Belgian Fonds de la Recherche Scientifique (FSR-FNRS). This work was supported by the Belgian Fonds de la Recherche Scientifique Médicale (FRSM).

> This poster will be made available for download after the meeting: http://www.facm.ucl.ac.be/posters.html

# A) Influence of acidic p A.1. Accumulation of a



# Acid pH markedly inc

B.1. Intrinsic activities of quinolones

Compounds -	MIC (mg/L)		
oompounds -	pH 7.4	pH 5.5	
Moxifloxacin	0.030	0.125	
Delafloxacin	0.0075	0.0002	

Acidic pH increases the activity of delafloxacin towards both extracellular and intracellular S. aureus, but impairs the activity of moxifloxacin.

Delafloxacin is highly intrinsically active against S. aureus. In contrast to moxifloxacin, it shows enhanced potency and efficacy at acidic pH against both extracellular and intracellular bacteria, in relation with its higher accumulation in both procaryotic and eucaryotic cells. This may confer an advantage to delafloxacin for the treatment of staphylococcal infections in territories where pH is acidic, including those involving intracellular reservoirs.

RESUL	TS				
oH on the accumulation of moxifloxacin and delafloxacin f antibiotics inside bacteria (S. <i>aureus</i> )	A.2. Accumulation				
Accumulation in <i>S. aureus:</i> Growing bacteria were incubated for 30 min in pH-adjust- ed broth (pH 7.4 vs. 5.5) with moxifloxacin (50 mg/L) or delafloxacin (100 mg/L). Values are means ± SD. Results are expressed as the cellu- lar to extracellular concentration ratio (% of control value [pH 7.4]).	Eluoroduinolone accumulation 20- 20- 20- 10- 10- 5- 0 Moxifloxa				
reases the accumulation of delafloxacin in both bacteria (S. aureus) or THP-					

# B) Influence of acidic pH on the activity of moxifloxacin and delafloxacin

B.2. Extracellular activities of quinolones



# CONCLUSIONS





UCL 73.70 av. Mounier 73, 1200 Brussels,

# on of antibiotics inside THP-1 cells



#### Accumulation in THP-1 cells:

THP-1 macrophages were incubated for 30 min. in cell culture medium (pH 7.4 vs. 5.5) with moxifloxacin (20 mg/L) or delafloxacin (20 mg/L). Values are means  $\pm$  SD. Results are expressed as the

cellular to extracellular concentration ratio (Cc/Ce).

# **1** cells. Opposite effects are observed for moxifloxacin.

# B.3. Intraphagocytic activities of quinolones

Extracellular (broth) and intracellular (THP-1 cells) activity of quinolones: Change in bacterial counts from the initial inoculum in broth (B2) or in infected cells (B3) after 24 h incubation in medium at pH 7.4 vs. pH 5.5 containing fixed extracellular concentrations of quinolones. All values are means ± SD of 3 independent determinations. Negative values denote a gain, and positive values, a loss in activity. Limit of detection: -5 log CFU.